

ROOT SYSTEMS AND WEYL GROUPOIDS FOR NICHOLS ALGEBRAS

I. HECKENBERGER AND H.-J. SCHNEIDER

ABSTRACT. Motivated by work of Kac and Lusztig, we define a root system and a Weyl groupoid for a large class of semisimple Yetter-Drinfeld modules over an arbitrary Hopf algebra. The obtained combinatorial structure fits perfectly into an existing framework of generalized root systems associated to a family of Cartan matrices, and provides novel insight into Nichols algebras. We demonstrate the power of our construction with new results on Nichols algebras over finite non-abelian simple groups and symmetric groups.

INTRODUCTION

In [Kac90], Kac defines the Lie algebra $\mathfrak{g}(A)$ associated to a symmetrizable Cartan matrix $A = (a_{ij})_{1 \leq i, j \leq n}$ by generators and relations, where the relations are not given explicitly, but they are determined by dividing out an ideal with a certain universal property. Similarly, Lusztig [Lus93] defines the braided algebra $\mathbf{f}$, that is the plus part of the quantum deformation of the the universal enveloping algebra of $\mathfrak{g}(A)$, by dividing out from the free algebra an ideal defined by a universal property (the radical of a bilinear form). After a considerable amount of work the relations turn out to be the Serre relations.

Using the language of braided vector spaces and braided categories, Lusztig's definition can be formulated as follows. Let V be a vector space with basis $x_1, \dots, x_n$ and define a braiding

$$c : V \otimes V \rightarrow V \otimes V, \quad c(x_i \otimes x_j) = q^{d_i a_{ij}} x_j \otimes x_i \text{ for all } i, j,$$

2000 *Mathematics Subject Classification.* 17B37, 16W30; 20F55.

Key words and phrases. Hopf algebra, quantum group, root system, Weyl group.

The work of I.H. was supported by DFG within a Heisenberg fellowship at the University of Munich.

where q is the deformation parameter and $(d_i a_{ij})$ is the symmetrized Cartan matrix. The braiding c has a categorical explanation. Let G be the free abelian group with basis $K_1, \dots, K_n$ and let ${}^G\mathcal{YD}$ be the category of Yetter-Drinfeld modules over G , that is, of G -graded vector spaces which are G -modules where each G -homogeneous part is stable under the action of G . Then $V \in {}^G\mathcal{YD}$, where each x_i has degree K_i and where the action is given by $K_i \cdot x_j = q^{d_i a_{ij}} x_j$ for all i, j , and V is a braided vector space as an object of the braided category ${}^G\mathcal{YD}$. Then

$$\mathbf{f} = T(V)/I_V,$$

where the free algebra $T(V)$ is a braided Hopf algebra such that the elements of V are primitive, and I_V is the largest coideal spanned by elements of $\mathbb{N}$ -degree ≥ 2 . In later terminology, $\mathfrak{B}(V) = T(V)/I_V$ is called the *Nichols algebra of V* . Note that $V = \mathbb{k}x_1 \oplus \dots \oplus \mathbb{k}x_n$ is a direct sum of one-dimensional Yetter-Drinfeld modules $\mathbb{k}x_i$, and the Nichols algebra of the irreducible pieces $\mathbb{k}x_i$ is easy to compute as a (truncated) commutative polynomial ring.

Yetter-Drinfeld modules can be defined over any Hopf algebra H over a field $\mathbb{k}$ with bijective antipode instead of the group algebra of G . For details we refer to Sect. 1. It is a fundamental problem in Hopf algebra theory to understand $\mathfrak{B}(V)$ for arbitrary objects V in the category ${}^H\mathcal{YD}$ of Yetter-Drinfeld modules over H (see the survey article [AS02b] of Andruskiewitsch and the second author). In this paper we study the Nichols algebra

$$\mathfrak{B}(M_1 \oplus \dots \oplus M_\theta), \quad \theta \geq 1, \quad M_1, \dots, M_\theta \in {}^H\mathcal{YD}$$

of the direct sum of finitely many finite-dimensional irreducible Yetter-Drinfeld modules $M_1, \dots, M_\theta$. We assume that finite tensor powers of $M_1 \oplus \dots \oplus M_\theta$ are semisimple in the category ${}^H\mathcal{YD}$ of Yetter-Drinfeld modules. Moreover, the braided adjoint action should satisfy a weak finiteness condition (see Def. 6.4). By Prop. 6.5, the finiteness condition can be checked directly using the braiding of $M_1 \oplus \dots \oplus M_\theta$. It holds for Lusztig's algebra $\mathbf{f}$. Our main achievement (Thms. 6.10, 6.11) is to associate to $M_1, \dots, M_\theta$ a generalized root system $\mathcal{R}$ in the sense of the paper [HY08] by Yamane and the first author. This is done similarly

to the definition of the root system of a Kac-Moody Lie algebra $\mathfrak{g}(A)$ in [Kac90]. Instead of one Cartan matrix A we have to deal with a family of Cartan matrices and with the Weyl groupoid of $\mathcal{R}$ instead of the Weyl group. In contrast to Lusztig's approach, the Cartan matrices are not given a priori, but they are obtained from the finiteness condition on the braided adjoint action.

The effect for the Nichols algebra is tremendous: Its dimension and (multivariate) Hilbert series are controlled by the set of roots, the Nichols algebras of $M_1, \dots, M_\theta$, and their images under reflections. The latter are much easier to calculate than the Nichols algebra itself. As for Kac-Moody Lie algebras, the situation is best if all roots are real. This is the case when the Weyl groupoid is finite, in particular when the Nichols algebra is finite-dimensional. Then we show in Thm. 7.2 the following striking result on the braided adjoint action: For all $i \neq j$ and $m \geq 1$, the Yetter-Drinfeld module $(\mathrm{ad}_c M_i)^m(M_j)$ is zero or irreducible.

To define the root system $\mathcal{R}$, we prove in Thm. 4.5 one of the main results in this paper: There exists a totally ordered index set $(L, \leq)$ and a family $(W_l)_{l \in L}$ of finite-dimensional irreducible $\mathbb{N}_0^\theta$ -graded objects in ${}^H_H\mathcal{YD}$ such that

$$\mathfrak{B}(M_1 \oplus \dots \oplus M_\theta) \simeq \otimes_{l \in L} \mathfrak{B}(W_l)$$

as $\mathbb{N}_0^\theta$ -graded objects in ${}^H_H\mathcal{YD}$, where $\deg N_i = \alpha_i$ for $1 \leq i \leq \theta$, and $\{\alpha_1, \dots, \alpha_\theta\}$ is the standard basis of $\mathbb{Z}^\theta$. Moreover, the Yetter-Drinfeld modules W_l and their degrees in $\mathbb{Z}^\theta$ are uniquely determined. The main ingredients of the proof are a generalization by Graña and the first author [GH07] of Kharchenko's result on PBW-bases of braided Hopf algebras of diagonal type [Kha99], and Thm. 3.5, which says that certain braided Hopf algebras are isomorphic as a Yetter-Drinfeld module to a Nichols algebra.

The second key part of the construction of our root system is the definition of the reflections and the proof of their basic properties. This is the main result of the paper [AHS08] of Andruskiewitsch and the authors.

Once the existence of the root system is established, we can use results on abstract generalized root systems recalled in Sect. 5. For recent work in this area we refer to [HY08] and the paper [CH08] by Cuntz and the first author.

Nichols algebras appear naturally inside of the associated graded Hopf algebra of pointed Hopf algebras [AS98], [AS02b]. Recall that a Hopf algebra is pointed if its simple comodules are one-dimensional. In particular, Hopf algebras generated by skew-primitive and group-like elements (as the quantum groups $U_q(\mathfrak{g}(A))$) are pointed. From our theory, we expect a deep impact on the further analysis of pointed Hopf algebras.

Our root system generalizes the construction of the first author [Hec06b] in the case of diagonal braidings, where H is the group algebra of an abelian group over a field of characteristic zero. In this setting, finite-dimensional Nichols algebras have been classified in [Hec06a]. These results allowed to complete the classification of a large class of finite-dimensional pointed Hopf algebras with abelian group of group-like elements [AS05].

The classification of finite-dimensional pointed Hopf algebras with non-abelian group of group-like elements is not known, but see [AZ07], [AF08] and the references therein. In the few examples, the group seems to be close to be abelian. This observation is supported by our results in Sect. 8, where the field is assumed to be algebraically closed of characteristic zero. In particular, in Cors. 8.3, 8.4 we show that the Nichols algebra of a non-simple Yetter-Drinfeld module over any non-abelian simple group or over the symmetric group $\mathbb{S}_n$, $n \geq 3$, is infinite dimensional. For arbitrary finite groups G we prove in Thm. 8.6 the following necessary condition for finiteness of the dimension of the Nichols algebra: If $i \neq j$ and if s respectively t are the degrees of nonzero homogeneous elements in M_i respectively M_j , then

$$(st)^2 = (ts)^2 \quad \text{in } G.$$

We believe that these rather immediate consequences of the existence of the root system form just the tip of the iceberg.

We would like to express our thanks to G. Malle for providing us with information on commuting conjugacy classes of finite groups.

1. PRELIMINARIES

Let $\mathbb{k}$ be a field and let H be a Hopf algebra over $\mathbb{k}$ with bijective antipode S . Recall that a (left) Yetter-Drinfeld module over H [Mon93, §10.6], [AS02b] is an H -module V equipped with a left co-module structure $\delta : V \rightarrow H \otimes V$, $v \mapsto v_{(-1)} \otimes v_{(0)}$, such that $\delta(h \cdot v) = h_{(1)}v_{(-1)}S(h_{(3)}) \otimes h_{(2)} \cdot v_{(0)}$ for all $h \in H$, $v \in V$.

Let ${}^H_H\mathcal{YD}$ and ${}^H_H\mathcal{YD}^{\text{fd}}$ denote the category of Yetter-Drinfeld modules and finite-dimensional Yetter-Drinfeld modules over H , respectively. The categories ${}^H_H\mathcal{YD}$ and ${}^H_H\mathcal{YD}^{\text{fd}}$ are braided with braiding

$$c : X \otimes Y \rightarrow Y \otimes X, \quad c(x \otimes y) = x_{(-1)} \cdot y \otimes x_{(0)}$$

for all $X, Y \in {}^H_H\mathcal{YD}$, $x \in X$, $y \in Y$. Braided bialgebras and Hopf algebras in this paper are always bialgebras and Hopf algebras in the braided category ${}^H_H\mathcal{YD}$.

If $B \in {}^H_H\mathcal{YD}$ is a braided Hopf algebra, $x, y \in B$, and x is a primitive element, then let $(\text{ad}_c x)(y) = xy - (x_{(-1)} \cdot y)x_{(0)}$.

Recall that a braided bialgebra is *connected*, if its coradical is $\mathbb{k}1$. Let ${}^H_H\mathcal{H}$ denote the category of connected $\mathbb{N}_0$ -graded braided Hopf algebras in ${}^H_H\mathcal{YD}$, which are generated as an algebra by elements of degree 1. The morphisms of ${}^H_H\mathcal{H}$ should be the maps of $\mathbb{N}_0$ -graded braided Hopf algebras. For any $R \in {}^H_H\mathcal{H}$ and $n \in \mathbb{N}_0$ we write $R(n)$ for the homogeneous component of R of degree n . Let $P(R)$ be the set of primitive elements of $R \in {}^H_H\mathcal{H}$. Let ${}^H_H\mathcal{H}^{\text{fg}}$ be the full subcategory of ${}^H_H\mathcal{H}$ consisting of those braided Hopf algebras R , for which $\dim R(1) < \infty$.

Definition 1.1. Let $R \in {}^H_H\mathcal{H}$. The maximal coideal of R contained in $(R^+)^2 = \bigoplus_{n \geq 2} R(n)$ is denoted by $\mathcal{I}_R$. Let $\pi_R : R \rightarrow R/\mathcal{I}_R$ denote the canonical projection.

If $P(R) = R(1)$, then R is called the *Nichols algebra* of $R(1)$ [AS02b]. Then we write $R = \mathfrak{B}(V)$, where $V = R(1) \in {}^H_H\mathcal{YD}$.

Let $R \in {}^H_H\mathcal{H}$. Then $\mathcal{I}_R$ is an $\mathbb{N}_0$ -graded Hopf ideal of R , and

$$(1.1) \quad R/\mathcal{I}_R \simeq \mathfrak{B}(R(1)).$$

Let $V \in {}^H_H\mathcal{YD}$. The tensor algebra $T(V)$ is a braided Hopf algebra in ${}^H_H\mathcal{YD}$ such that $P(T(V)) = V$. Another description of the ideal $\mathfrak{I}_{T(V)}$ is as the sum of the kernels of the quantum symmetrizer $S_n \in \text{End}(V^{\otimes n})$, $n \geq 2$, introduced by Woronowicz [Wor89], see [Sch96], [Maj95, 10.4.13]. Explicitly, S_n can be defined with help of the quantum shuffle maps $S_{k,1} \in \text{End}(V^{\otimes k+1})$, $k \geq 1$, as follows. We write $c_{i,i+1}$, if we apply the braiding to the i -th and $i+1$ -st components of a tensor product of Yetter-Drinfeld modules. Then

$$(1.2) \quad S_{n-1,1} = \sum_{k=1}^n c_{n-1,n} c_{n-2,n-1} \cdots c_{k,k+1},$$

$$(1.3) \quad S_n = (S_{1,1} \otimes \text{id})(S_{2,1} \otimes \text{id}) \cdots (S_{n-2,1} \otimes \text{id}) S_{n-1,1}.$$

For any $\mathbb{Z}^n$ -graded vector space X , where $n \in \mathbb{N}$, let X_γ denote the homogeneous component of degree $\gamma \in \mathbb{Z}^\theta$ of X .

2. APPROXIMATIONS OF NICHOLS ALGEBRAS

In this section we define and study for each $i \in \mathbb{N} \cup \{\infty\}$ a functor $F_i : {}^H_H\mathcal{H} \rightarrow {}^H_H\mathcal{H}$ and a natural transformation λ_i from id to F_i . The functor F_i converts primitive elements of degree at most i to elements of degree 1, and λ_i maps primitive elements of degree 2, 3, $\dots$, i to zero.

Definition 2.1. Let $R = \bigoplus_{n=0}^\infty R(n) \in {}^H_H\mathcal{H}$ and $i \in \mathbb{N} \cup \{\infty\}$. Let

$$(2.1) \quad P_i(R) = \bigoplus_{n=1}^i R(n) \cap P(R),$$

$$(2.2) \quad R_{-1} = 0, \quad R_0 = \mathbb{k}, \quad R_k = (\mathbb{k} \oplus P_i(R))^k \subset R \quad \text{for all } k \in \mathbb{N},$$

$$(2.3) \quad F_i(R) = \bigoplus_{k=0}^\infty R_k / R_{k-1}.$$

Let $\lambda_i : R \rightarrow F_i(R)$ be the linear map defined by

$$(2.4) \quad \lambda_i : R(k) \ni x \mapsto x + R_{k-1} \in R_k / R_{k-1} \quad \text{for all } k \in \mathbb{N}_0.$$

For later use we define

$$(2.5) \quad P'_i(R) = \bigoplus_{n=2}^i R(n) \cap P(R).$$

Remark 2.2. Let $R = \bigoplus_{n=0}^\infty R(n) \in {}^H_H\mathcal{H}$ and $i \in \mathbb{N} \cup \{\infty\}$. Then $R_k / R_{k-1} \in {}^H_H\mathcal{YD}$ for all $k \in \mathbb{N}_0$, and $F_i(R) \in {}^H_H\mathcal{H}$ with algebra and coalgebra structure induced by those of R , and $\mathbb{N}_0$ -grading given by the decomposition in Eq. (2.3). If $i \in \mathbb{N}$ and $R \in {}^H_H\mathcal{H}^{\text{fg}}$, then $\dim R(k) < \infty$

for all $k \in \mathbb{N}$, since $R(k) = R(1)^k$. Therefore in this case $P_i(R)$ is finite-dimensional, and hence $F_i(R) \in {}^H_H\mathcal{H}^{\text{fg}}$.

Let $R, S \in {}^H_H\mathcal{H}$ and $f : R \rightarrow S$ a morphism. Then f induces a morphism $F_i(f) : F_i(R) \rightarrow F_i(S)$ in ${}^H_H\mathcal{H}$. Thus F_i is a covariant functor.

The map $\lambda_i : R \rightarrow F_i(R)$ is well-defined, since $R(k) = R(1)^k$ and $R(1) \subset P_i(R)$, and it is a morphism in ${}^H_H\mathcal{H}$. Moreover, $\ker \lambda_i$ is the ideal of R generated by $P'_i(R)$, and $\text{Im } \lambda_i$ is the subalgebra of $F_i(R)$ generated by $R(1) \subset F_i(R)(1)$.

Proposition 2.3. *Let $R \in {}^H_H\mathcal{H}$ and $i \in \mathbb{N} \cup \{\infty\}$. Let $\mathbb{k}\langle R(1) \rangle$ be the subalgebra of $F_i(R)$ generated by $R(1) \subset F_i(R)(1)$, and let $(P'_i(R))$ be the ideal of $F_i(R)$ generated by $P'_i(R) \subset F_i(R)(1)$. Then $\mathbb{k}\langle R(1) \rangle \in {}^H_H\mathcal{H}$, $(P'_i(R))$ is an $\mathbb{N}_0$ -graded Hopf ideal of $F_i(R)$ in ${}^H_H\mathcal{YD}$, and $F_i(R) = \mathbb{k}\langle R(1) \rangle \oplus (P'_i(R))$.*

Proof. Clearly, $\mathbb{k} \oplus R(1)$ is a subcoalgebra and Yetter-Drinfeld submodule of $F_i(R)$, and hence $\mathbb{k}\langle R(1) \rangle \in {}^H_H\mathcal{H}$. Further, $(P'_i(R)) \in {}^H_H\mathcal{YD}$ since $P'_i(R) \in {}^H_H\mathcal{YD}$, and $(P'_i(R))$ is an $\mathbb{N}_0$ -graded coideal of $F_i(R)$ since $P'_i(R)$ is an $\mathbb{N}_0$ -graded coideal of $F_i(R)$. The algebra $F_i(R)$ is generated by $R(1) \oplus P'_i(R)$, and therefore $\mathbb{k}\langle R(1) \rangle + (P'_i(R)) = F_i(R)$. It remains to show that the sum is direct. Since $\mathbb{k}\langle R(1) \rangle$ and $(P'_i(R))$ are $\mathbb{N}_0$ -graded, it suffices to consider $\mathbb{N}_0$ -homogeneous components. Let $k \geq 0$ and $\bar{x} \in \mathbb{k}\langle R(1) \rangle \cap (P'_i(R)) \cap F_i(R)(k)$. Since $\bar{x} \in \mathbb{k}\langle R(1) \rangle$, there exists a representant $x \in R(k)$ of $\bar{x} \in R_k/R_{k-1}$. Further, $\bar{x} \in (P'_i(R))$, and hence the set $x + R_{k-1}$ contains an element y of the ideal $(P'_i(R)) \subset R$. By subtracting elements of R_{k-1} from y we may assume that y is a linear combination of monomials containing k factors from $R(1) \cup P'_i(R)$, and in each monomial at least one of the factors is from $P'_i(R)$. But $P'_i(R) \subset \bigoplus_{j \geq 2} R(j)$, and hence $y \in \bigoplus_{j \geq k+1} R(j)$. Since $x \in R(k)$ and $x - y \in R_{k-1}$, this implies that $x = y = 0$ in R_k/R_{k-1} , and hence $\mathbb{k}\langle R(1) \rangle \cap (P'_i(R)) = 0$. $\square$

Remark 2.4. Let $R \in {}^H_H\mathcal{H}$ and let $R(1) = V \oplus W$ be a decomposition in ${}^H_H\mathcal{YD}$. Let $\mathbb{k}\langle V \rangle$ be the subalgebra of R generated by V , and let (W) be the ideal of R generated by W . Then in general the sum $R = \mathbb{k}\langle V \rangle + (W)$ is not direct.

For example, let $H = \mathbb{k}(\mathbb{Z}/2\mathbb{Z})^2$ and assume that $g, h \in (\mathbb{Z}/2\mathbb{Z})^2$ generate $(\mathbb{Z}/2\mathbb{Z})^2$ as a group. Let $V, W \in {}^H_H\mathcal{YD}$ with $V = \mathbb{k}x$, $W = \mathbb{k}y$, and assume that the coaction and the action of H satisfy

$$\begin{aligned}\delta(x) &= g \otimes x, & \delta(y) &= h \otimes y, \\ g \cdot x &= -x, & g \cdot y &= y, & h \cdot x &= x, & h \cdot y &= -y.\end{aligned}$$

Then $\mathfrak{J} = (xy - yx, x^2 - y^2)$ is an $\mathbb{N}_0$ -graded Hopf ideal in $T(V \oplus W)$, where $V \oplus W \subset P(T(V \oplus W))$. Since $x^2 \in \mathbb{k}\langle V \rangle \cap (W) \subset T(V \oplus W)/\mathfrak{J}$, the sum $\mathbb{k}\langle V \rangle + (W)$ in $T(V \oplus W)/\mathfrak{J}$ is not direct.

Lemma 2.5. *Let $R \in {}^H_H\mathcal{H}$, $k \in \mathbb{N}_0$, and $i \in \mathbb{N}$ such that $i > k$. If $R(m) \cap P(R) = 0$ for all m with $2 \leq m \leq k$, then $\lambda_i(R)(m) \cap P(\lambda_i(R)) = 0$ for all m with $2 \leq m \leq k + 1$.*

Proof. The condition $R(m) \cap P(R) = 0$ for all m with $2 \leq m \leq k$ is equivalent to $P'_i(R) = \bigoplus_{n=k+1}^i R(n) \cap P(R)$. Since $\lambda_i(R) \simeq R/\ker \lambda_i$, the claim follows from the end of Rem. 2.2. $\square$

Proposition 2.6. *Let $m \in \mathbb{N}$ and $i_2, i_3, \dots, i_m \in \mathbb{N}$ such that $i_n \geq n$ for all $n \leq m$. Let $R \in {}^H_H\mathcal{H}$ and*

$$\lambda = \lambda_{i_m} \cdots \lambda_{i_3} \lambda_{i_2} : R \rightarrow F_{i_m} \cdots F_{i_3} F_{i_2}(R).$$

Then $\ker \lambda \cap R(k) = \mathfrak{J}_R \cap R(k)$ for all k with $0 \leq k \leq m$.

Proof. By Lemma 2.5 and induction on m one obtains that $\lambda(R(k))$ has, if $2 \leq k \leq m$, no primitive elements besides 0. Since R is generated by $R(1)$ and $\lambda(R) \simeq R/\ker \lambda$, we obtain that $\ker \lambda \cap R(k) = \mathfrak{J}_R \cap R(k)$ for $0 \leq k \leq m$. Now, if $i \leq m$, then $P'_i(F_{i_m} \cdots F_{i_3} F_{i_2}(R)) \cap \lambda(R) = 0$, and hence $\lambda_i : \lambda(R) \rightarrow \lambda_i \lambda(R)$ is an isomorphism. Otherwise $i > m$, and the claim follows from the equalities

$$\ker \lambda \cap R(k) = \ker \lambda_i \lambda \cap R(k) = \mathfrak{J}_R \cap R(k), \quad 0 \leq k \leq m,$$

shown in the first part of the proof. $\square$

3. BRAIDED HOPF ALGEBRAS AND YETTER-DRINFELD MODULES

In Thm. 3.5 we give a criterion for a braided Hopf algebra to be isomorphic to a Nichols algebra in ${}^H_H\mathcal{YD}$. First we show that $F_i(R) \simeq R$ in ${}^H_H\mathcal{YD}$ for all $R \in {}^H_H\mathcal{H}$ which are semisimple objects in ${}^H_H\mathcal{YD}$.

Lemma 3.1. *Let R be a coalgebra in ${}^H_H\mathcal{YD}$ and let $R = \cup_{n=0}^{\infty} R_n$ be a coalgebra filtration of R in ${}^H_H\mathcal{YD}$. Suppose that R_n is a semisimple object in ${}^H_H\mathcal{YD}$ for all $n \in \mathbb{N}_0$. Let $\text{gr } R = \oplus_{n=0}^{\infty} R(n)$, where $R_{-1} = 0$ and $R(n) = R_n/R_{n-1}$ for all $n \in \mathbb{N}_0$. Then the coalgebra $\text{gr } R$ is a semisimple object in ${}^H_H\mathcal{YD}$, and $\text{gr } R \simeq R$ as objects in ${}^H_H\mathcal{YD}$.*

Proof. For each $n \in \mathbb{N}_0$ the sequence

$$0 \rightarrow R_{n-1} \rightarrow R_n \rightarrow R(n) \rightarrow 0$$

in ${}^H_H\mathcal{YD}$ is exact. Thus $R_n \simeq R(n) \oplus R_{n-1}$ in ${}^H_H\mathcal{YD}$ and $R(n)$ is semisimple for all $n \in \mathbb{N}_0$, since R_n is semisimple for all $n \in \mathbb{N}_0$. By induction one obtains for all $n \in \mathbb{N}_0$ that $R_n \simeq \oplus_{i=0}^n R(i)$ in ${}^H_H\mathcal{YD}$. For all $n \in \mathbb{N}_0$ the isomorphism $R_n \simeq \oplus_{i=0}^n R(i)$ can be extended to an isomorphism $R_{n+1} \simeq \oplus_{i=0}^{n+1} R(i)$, since $R_{n+1} \simeq R_n \oplus R(n+1)$. Thus the lemma follows from

$$R = \cup_{n \in \mathbb{N}} R_n \simeq \cup_{n \in \mathbb{N}} \oplus_{i=0}^n R(i) \simeq \oplus_{i=0}^{\infty} R(i) = \text{gr } R. \quad \square$$

Lemma 3.2. *Let R be an algebra in ${}^H_H\mathcal{YD}$ which is generated by a subspace $U \in {}^H_H\mathcal{YD}$. Let $R_{-1} = 0$, $R_0 = \mathbb{k}$, $R_n = (U + \mathbb{k})^n \subset R$ for all $n \in \mathbb{N}$, and $\text{gr } R = \oplus_{n=0}^{\infty} R_n/R_{n-1}$. If R_n is semisimple in ${}^H_H\mathcal{YD}$ for all $n \in \mathbb{N}_0$, then $\text{gr } R \simeq R$ as objects in ${}^H_H\mathcal{YD}$.*

Proof. Since $\mathbb{k} \otimes W \simeq W$ for all $W \in {}^H_H\mathcal{YD}$, R_{n-1} is a subobject of R_n in ${}^H_H\mathcal{YD}$ for all $n \in \mathbb{N}_0$. Thus the arguments in the proof of Lemma 3.1 yield the claim. $\square$

Corollary 3.3. *Let $R \in {}^H_H\mathcal{H}$ and $i \in \mathbb{N}$. If $R(n)$ is semisimple in ${}^H_H\mathcal{YD}$ for all $n \in \mathbb{N}$, then $F_i(R)(n)$ is semisimple in ${}^H_H\mathcal{YD}$ for all $n \in \mathbb{N}$, and $F_i(R) \simeq R$ as objects in ${}^H_H\mathcal{YD}$.*

Proof. This is clear by the definition of F_i and by Lemma 3.2. $\square$

Recall that for any embedding $\iota_U^V : U \subset V$ in ${}^H_H\mathcal{YD}$ there exists a canonical embedding $\mathfrak{B}(\iota_U^V) : \mathfrak{B}(U) \rightarrow \mathfrak{B}(V)$ of braided Hopf algebras in ${}^H_H\mathcal{YD}$ such that $\mathfrak{B}(\iota_U^V)|_U = \iota_U^V$.

Lemma 3.4. *Let $R \in {}^H_H\mathcal{H}$ and $i \in \mathbb{N}$ such that all finite tensor powers of $R(1)$ are semisimple in ${}^H_H\mathcal{YD}$. Let $\iota_R : \mathfrak{B}(R(1)) \rightarrow R$ be an $\mathbb{N}_0$ -graded*

splitting of the exact sequence

$$0 \rightarrow \mathfrak{I}_R \longrightarrow R \xrightarrow{\pi_R} \mathfrak{B}(R(1)) \rightarrow 0$$

in ${}^H_H\mathcal{YD}$. Then there is an isomorphism $\Phi : R \rightarrow F_i(R)$ in ${}^H_H\mathcal{YD}$ and an $\mathbb{N}_0$ -graded splitting $\iota_{F_i(R)} : \mathfrak{B}(P_i(R)) \rightarrow F_i(R)$ of the exact sequence

$$0 \rightarrow \mathfrak{I}_{F_i(R)} \longrightarrow F_i(R) \xrightarrow{\pi_{F_i(R)}} \mathfrak{B}(P_i(R)) \rightarrow 0$$

in ${}^H_H\mathcal{YD}$ with the following properties.

- (1) $\Phi(R(n)) \subset \oplus_{k=0}^n F_i(R)(k)$ for all $n \in \mathbb{N}_0$.
- (2) Let $\text{gr } \Phi \in \text{Hom}(R, F_i(R))$ such that $\text{gr } \Phi|_{R(n)} = \text{pr}_n \circ \Phi|_{R(n)}$ for all $n \in \mathbb{N}_0$, where $\text{pr}_n : F_i(R) \rightarrow F_i(R)(n)$ is the canonical projection. Then $\text{gr } \Phi = \lambda_i$.
- (3) The following diagram is commutative.

$$(3.1) \quad \begin{array}{ccc} \mathfrak{B}(R(1)) & \xrightarrow{\iota_R} & R \\ \mathfrak{B}(\iota_{R(1)}^{P_i(R)}) \downarrow & & \downarrow \Phi \\ \mathfrak{B}(P_i(R)) & \xrightarrow{\iota_{F_i(R)}} & F_i(R). \end{array}$$

Proof. First we define the map $\Phi|_{R(k)}$ for all $k \in \mathbb{N}_0$ and prove that it satisfies (1) and (2). Then we show that Φ is bijective, and that the required section $\iota_{F_i(R)}$ of $\pi_{F_i(R)}$, which makes Diagram (3.1) commutative, exists.

For all $k \in \mathbb{N}_0$ let $(P'_i(R))(k) = (P'_i(R)) \cap R(k)$, where $(P'_i(R)) \subset R$ is the ideal of R generated by $P'_i(R)$. Since $P'_i(R) \subset \oplus_{n=2}^{\infty} R(n)$ is an $\mathbb{N}_0$ -graded object in ${}^H_H\mathcal{YD}$, we obtain that $(P'_i(R)) = \oplus_{k=2}^{\infty} (P'_i(R))(k)$ and that $(P'_i(R))(k) \in {}^H_H\mathcal{YD}$ for all $k \in \mathbb{N}_0$. Further, the sum

$$(3.2) \quad \iota_R(\mathfrak{B}^k(R(1))) + (P'_i(R))(k) \subset R(k)$$

is direct. Indeed, π_R is an algebra map, $\pi_R(P'_i(R)) = 0$, and $\pi_R \iota_R = \text{id}_{\mathfrak{B}(R(1))}$. Hence, if $x \in \mathfrak{B}^k(R(1))$ and $\iota_R(x) \in (P'_i(R))(k)$, then $\pi_R(x) \in \pi_R((P'_i(R))) = 0$ and $\pi_R(x) = \pi_R \iota_R(x) = x$, and hence $x = 0$.

For all $j, k \in \mathbb{N}$ let $V_j^{ik} \in {}^H_H\mathcal{YD}$ such that

$$R(k) = (R_j \cap R(k)) \oplus V_j^{ik}$$

(see Eq. (2.2)). The V_j^{ik} exist, since $R(k)$ is semisimple in ${}^H_H\mathcal{YD}$. The relation $P'_i(R) \subset \oplus_{n=2}^\infty R(n)$ implies that

$$(P'_i(R))(k) = R_{k-1} \cap (P'_i(R))(k) = R_{k-1} \cap R(k),$$

and hence by Eq. (3.2) we may assume that $\iota_R(\mathfrak{B}^k(R(1))) \subset V_j^{ik}$ if $j < k$. Let $k \in \mathbb{N}_0$. We define $\Phi_k : R(k) \rightarrow F_i(R)$ by setting

$$(3.3) \quad \begin{aligned} \Phi_k &:= \oplus_{j=0}^k \Phi_{kj}, & \Phi_{kj} : R(k) &\rightarrow F_i(R)(j) = R_j/R_{j-1}, \\ \Phi_{kj}(x) &:= \begin{cases} 0 & \text{for } x \in V_j^{ik}, \\ x + R_{j-1} & \text{for } x \in R_j \cap R(k), \end{cases} \end{aligned}$$

where $0 \leq j \leq k$. Then it is clear that (1) holds, and since $\text{gr } \Phi|_{R(k)} = \Phi_{kk}$, we also get (2) by the definition of λ_i in Eq. (2.4).

Now we prove that Φ is bijective. Suppose first that $x = \sum_{k=0}^n x_k$, where $x_k \in R(k)$ for all $k \leq n$, and $\Phi(x) = 0$. Then

$$\Phi(x) = \sum_{j=0}^n \sum_{k=j}^n \Phi_{kj}(x_k) \in \sum_{j=0}^n F_i(R)(j)$$

by definition of Φ and by (1), and hence $\sum_{k=j}^n \Phi_{kj}(x_k) = 0$ for all $j \in \{0, 1, \dots, n\}$. Using the definition of Φ again and again, and the facts that $R(k) \subset R_k$ and R_k is $\mathbb{N}_0$ -graded for all $k \in \mathbb{N}_0$, we conclude now as follows.

$$\begin{aligned} \Phi_{nn}(x_n) = 0 &\Rightarrow x_n \in R_{n-1}, \\ \Phi_{n-1\,n-1}(x_{n-1}) + \Phi_{nn}(x_n) = 0 &\Rightarrow x_{n-1} + x_n \in R_{n-2} \Rightarrow \\ x_{n-1}, x_n &\in R_{n-2}, \\ \Phi_{n-2\,n-2}(x_{n-2}) + \Phi_{n-1\,n-2}(x_{n-1}) + \Phi_{nn}(x_n) = 0 &\Rightarrow \\ x_{n-2} + x_{n-1} + x_n &\in R_{n-3} \Rightarrow x_k \in R_{n-3} \text{ if } n-2 \leq k \leq n, \dots \\ \sum_{k=1}^n x_k \in R_0 = \mathbb{k} &\Rightarrow x_k = 0 \text{ if } 1 \leq k \leq n, \\ \Phi_{00}(x_0) = 0 &\Rightarrow x_0 \in R_{-1} = 0. \end{aligned}$$

Thus $x_k = 0$ for all $k \in \{0, 1, \dots, n\}$, and hence Φ is injective. The surjectivity of Φ follows immediately from the decomposition

$$\begin{aligned} R_j/R_{j-1} &= ((R(1) \oplus P'_i(R))^j + R_{j-1})/R_{j-1} \\ &= \bigoplus_{k=j}^{\infty} (R(k) \cap R_j + R_{j-1})/R_{j-1} \\ &= \bigoplus_{k=j}^{\infty} (R(k) \cap R_j + R(k) \cap R_{j-1})/(R(k) \cap R_{j-1}) \end{aligned}$$

and the definition of Φ .

Now we prove (3). Note that the canonical map $\pi_{F_i(R)} : F_i(R) \rightarrow \mathfrak{B}(P_i(R))$ is compatible with the decomposition $F_i(R) = \mathbb{k}\langle R(1) \rangle \oplus (P'_i(R))$, see Prop. 2.3, in the sense that $\pi_{F_i(R)}(\mathbb{k}\langle R(1) \rangle) \subset \mathbb{k}\langle R(1) \rangle$ and $\pi_{F_i(R)}((P'_i(R))) \subset (P'_i(R))$. Let $\pi' : \mathbb{k}\langle R(1) \rangle \rightarrow \mathbb{k}\langle R(1) \rangle$ and $\pi'' : (P'_i(R)) \rightarrow (P'_i(R))$ denote the components of $\pi_{F_i(R)}$, that is, $\pi_{F_i(R)} = \pi' \oplus \pi''$. Let $\iota''_{F_i(R)} : (P'_i(R)) \rightarrow (P'_i(R))$ be an arbitrary section of π'' in ${}^H_H\mathcal{YD}$, and define

$$\iota'_{F_i(R)} : \mathbb{k}\langle R(1) \rangle \rightarrow \mathbb{k}\langle R(1) \rangle, \quad \iota'_{F_i(R)} := \lambda_i \circ \iota_R,$$

where the domain of $\iota'_{F_i(R)}$ is a subobject of $\mathfrak{B}(P'_i(R))$, and the range of $\iota'_{F_i(R)}$ is a subobject of $F_i(R)$. Let $\iota_{F_i(R)} := \iota'_{F_i(R)} \oplus \iota''_{F_i(R)}$. Note that the choice of $\iota'_{F_i(R)}$ is necessary. Indeed, in order to make Diagram (3.1) commutative, we need precisely that $\iota'_{F_i(R)} = \Phi \circ \iota_R$. However, Eq. (3.3) and the assumption $\iota_R(\mathfrak{B}^k(R(1))) \subset V_j^{ik}$ for $j < k$ imply that $\Phi_{kj}\iota_R(\mathfrak{B}^k(R(1))) = 0$ for $j < k$, and hence $\Phi\iota_R = (\text{gr } \Phi)\iota_R = \lambda_i\iota_R$.

It remains to show that $\pi_{F_i(R)}\iota'_{F_i(R)}(\mathbb{k}\langle R(1) \rangle) = \text{id}|_{\mathbb{k}\langle R(1) \rangle}$. Equivalently, we have to prove that the diagram

$$\begin{array}{ccc} \mathfrak{B}(R(1)) & \xrightarrow{\iota_R} & R \\ \mathfrak{B}(\iota_{R(1)}^{P_i(R)}) \downarrow \simeq & & \downarrow \lambda_i \\ \mathfrak{B}(P_i(R)) \supset \mathbb{k}\langle R(1) \rangle & \xleftarrow{\pi_{F_i(R)}} & \mathbb{k}\langle R(1) \rangle \subset F_i(R) \end{array}$$

is commutative. Clearly, this is equivalent to the commutativity of the diagrams

$$\begin{array}{ccccccc} \mathfrak{B}(R(1)) & \xrightarrow[\simeq]{\iota_R} & \iota_R(\mathfrak{B}(R(1))) & & \mathfrak{B}(R(1)) & \xleftarrow[\simeq]{\pi_R} & \iota_R(\mathfrak{B}(R(1))) \\ \mathfrak{B}(\iota_{R(1)}^{P_i(R)}) \downarrow & & \downarrow \lambda_i & & \mathfrak{B}(\iota_{R(1)}^{P_i(R)}) \downarrow & & \downarrow \lambda_i \\ \mathfrak{B}(P_i(R)) & \xleftarrow{\pi_{F_i(R)}} & F_i(R), & & \mathfrak{B}(P_i(R)) & \xleftarrow{\pi_{F_i(R)}} & F_i(R), \end{array}$$

where the second one is obtained from the first one by using that ι_R is a section of π_R . The second diagram is easily seen to be commutative. Indeed, the diagram

$$\begin{array}{ccc} \mathfrak{B}(R(1)) & \xleftarrow{\pi_R} & R \\ \mathfrak{B}(\iota_{R(1)}^{P_i(R)}) \downarrow & & \downarrow \lambda_i \\ \mathfrak{B}(P_i(R)) & \xleftarrow{\pi_{F_i(R)}} & F_i(R), \end{array}$$

is a diagram in ${}^H_H\mathcal{H}$, and it is commutative, since it is commutative on the generators. $\square$

Theorem 3.5. *Let R be a connected braided Hopf algebra which is generated as an algebra by $U \subset P(R)$, where $U \in {}^H_H\mathcal{YD}$, and assume that all tensor powers of U are semisimple in ${}^H_H\mathcal{YD}$. Then there is a subobject $V \subset R$ in ${}^H_H\mathcal{YD}$ such that $U \subset V$, all tensor powers of V are semisimple in ${}^H_H\mathcal{YD}$, and $R \simeq \mathfrak{B}(V)$ in ${}^H_H\mathcal{YD}$.*

Proof. It is sufficient to prove the theorem for $R \in {}^H_H\mathcal{H}$ and $U = R(1)$. Indeed, define $\text{gr } R = \bigoplus_{n=0}^{\infty} R(n)$ as in Lemma 3.2. Since $U \subset P(R)$, we get $\Delta : R(n) \rightarrow \bigoplus_{k=0}^n R(k) \otimes R(n-k)$, and hence $\text{gr } R \in {}^H_H\mathcal{H}$. Moreover, $\text{gr } R \simeq R$ in ${}^H_H\mathcal{YD}$ by Lemma 3.2.

Define inductively $R^{(n)} \in {}^H_H\mathcal{H}$ for all $n \in \mathbb{N}$ by letting $R^{(1)} = R$ and $R^{(n)} = F_{n-1}(R^{(n-1)})$ for all $n \in \mathbb{N}_{\geq 2}$, and let $P_n = P_n(R^{(n)})$, $P'_n = P'_n(R^{(n)})$ for all $n \in \mathbb{N}$.

Using Corollary 3.3 and the formula $R^{(n)}(k) = R^{(n)}(1)^k$ for all $k, n \in \mathbb{N}$, we may apply Lemma 3.4. Hence for each $n \in \mathbb{N}$ there exists an isomorphism $\Phi_n : R^{(n)} \rightarrow R^{(n+1)}$ which satisfies the properties in Lemma 3.4(1)–(3). In particular, for each $n \in \mathbb{N}_{\geq 2}$ the diagrams

$$(3.4) \quad \begin{array}{ccccc} \mathfrak{B}(P_{n-1}) & \xrightarrow{\iota_{R^{(n)}}} & R^{(n)} & \mathfrak{B}(P_{n-1}) & \xrightarrow{\Phi_1^{-1} \dots \Phi_{n-1}^{-1} \iota_{R^{(n)}}} & R \\ \mathfrak{B}(\iota_{P_{n-1}}^{P_n}) \downarrow & & \downarrow \Phi_n & \mathfrak{B}(\iota_{P_{n-1}}^{P_n}) \downarrow & & \parallel \\ \mathfrak{B}(P_n) & \xrightarrow{\iota_{R^{(n+1)}}} & R^{(n+1)}, & \mathfrak{B}(P_n) & \xrightarrow{\Phi_1^{-1} \dots \Phi_n^{-1} \iota_{R^{(n+1)}}} & R \end{array}$$

commute. Hence there is a map

$$\Psi : \mathfrak{B}(\bigcup_{n=1}^{\infty} P_n) = \bigcup_{n=1}^{\infty} \mathfrak{B}(P_n) \rightarrow R$$

in ${}^H_H\mathcal{YD}$, and since $\iota_{R^{(n)}}$ is injective for all $n \in \mathbb{N}$, the map Ψ is injective. Let $V = \cup_{n=0}^\infty P_n \in {}^H_H\mathcal{YD}$. Since for all $n \in \mathbb{N}$ all tensor powers of P_n are semisimple in ${}^H_H\mathcal{YD}$, all tensor powers of V are semisimple in ${}^H_H\mathcal{YD}$. We are left to show that Ψ is surjective.

For all $n \in \mathbb{N}$ let $\varphi_n = \Phi_{n-1} \cdots \Phi_2 : R^{(2)} = R \rightarrow R^{(n)}$. By Lemma 3.4, for all $x \in R$ there exist $k \in \mathbb{N}_0$ and $m \in \mathbb{N}_{\geq 2}$ such that

$$\varphi_m(x) \in \oplus_{n=0}^k R^{(m)}(n), \quad \varphi_t(x) \notin \oplus_{n=0}^{k-1} R^{(t)}(n)$$

for all $t \geq m$. We prove by induction on k that $x \in \Psi(\mathfrak{B}(V))$. First, let $k = 0$. Then $x \in \mathbb{k} = \mathfrak{B}^0(V)$, and $\Psi(x) = x$ by definition of Ψ .

Let now $k \geq 1$. Since

$$\varphi_m(x) - \iota_{R^{(m)}} \pi_{R^{(m)}} \varphi_m(x) \in R^{(m)}(k) \cap \ker \pi_{R^{(m)}} = R^{(m)}(k) \cap \mathfrak{I}_{R^{(m)}},$$

Prop. 2.6 and Lemma 3.4(2) imply that

$$\Phi_{m+k-2} \cdots \Phi_{m+1} \Phi_m(\varphi_m(x) - \iota_{R^{(m)}} \pi_{R^{(m)}} \varphi_m(x)) \in \oplus_{n=0}^{k-1} R^{(m+k-1)}(n).$$

Now $\pi_{R^{(m)}} \varphi_m(x) \in \mathfrak{B}(V)$ by definition of $\mathfrak{B}(V)$, and hence

$$\varphi_m^{-1} \iota_{R^{(m)}} \pi_{R^{(m)}} \varphi_m(x) \in \Psi(\mathfrak{B}(V)).$$

Further, $x - \varphi_m^{-1} \iota_{R^{(m)}} \pi_{R^{(m)}} \varphi_m(x) \in \Psi(\mathfrak{B}(V))$ by induction hypothesis on k . Thus $x \in \Psi(\mathfrak{B}(V))$. $\square$

Remark 3.6. Assume that $R \in {}^H_H\mathcal{YD}^{\text{fd}}$ is a connected braided Hopf algebra and a semisimple object in ${}^H_H\mathcal{YD}^{\text{fd}}$. Then, using the algorithm in [AS02a] below Cor. 7.8, it can be shown that $R \simeq \mathfrak{B}(V)$ in ${}^H_H\mathcal{YD}^{\text{fd}}$ for some $V \in {}^H_H\mathcal{YD}^{\text{fd}}$.

4. DECOMPOSITIONS OF NICHOLS ALGEBRAS INTO TENSOR PRODUCTS

Our main result in this section is Thm. 4.5 giving a decomposition of a large class of Nichols algebras. For the proof we use Thm. 3.5 and a result from [GH07] which is based on the ideas of Kharchenko on PBW bases of braided Hopf algebras with diagonal braiding [Kha99].

Let ${}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$ denote the category of $\mathbb{Z}^\theta$ -graded Yetter-Drinfeld modules over H having finite-dimensional homogeneous components.

Remark 4.1. Let H' denote the Hopf algebra $\mathbb{k}\mathbb{Z}^\theta \otimes H$. For any $\mathbb{Z}^\theta$ -graded object $V \in {}^H_H\mathcal{YD}$ let $V' \in {}^{H'}_{H'}\mathcal{YD}$ such that $V' = V$ as an H -module, $\gamma v = v$ for all $v \in V'$ and $\gamma \in \mathbb{Z}^\theta$, and the left coaction on V' is determined by $\delta_{V'}(v) = \gamma v_{(-1)} \otimes v_{(0)}$ for all $v \in V_\gamma \subset V'$, where $v_{(-1)} \otimes v_{(0)}$ is the coaction of H on $v \in V$. This way the category of $\mathbb{Z}^\theta$ -graded objects in ${}^H_H\mathcal{YD}$ is equivalent to the full subcategory of ${}^{H'}_{H'}\mathcal{YD}$ consisting of those Yetter-Drinfeld modules, for which the action of $\mathbb{k}\mathbb{Z}^\theta$ is trivial.

Let $\mathcal{A} = \{\alpha_1, \dots, \alpha_\theta\}$ be a fixed basis of $\mathbb{Z}^\theta$. For $\gamma = \sum_{i=1}^\theta n_i \alpha_i \in \mathbb{Z}^\theta$ let $|\gamma|_{\mathcal{A}} = \sum_{i=1}^\theta n_i$. For any $\mathbb{Z}^\theta$ -graded object X and any $k \in \mathbb{Z}$ let

$$X(k) = \bigoplus_{\gamma \in \mathbb{Z}^\theta, |\gamma|_{\mathcal{A}}=k} X_\gamma.$$

Clearly, if $X \in {}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$ such that $X_\gamma = 0$ for $\gamma \in \mathbb{Z}^\theta \setminus \mathbb{N}_0\mathcal{A}$, where $\mathbb{N}_0\mathcal{A} = \sum_{i=1}^\theta \mathbb{N}_0\alpha_i$, then $X(k) = 0$ for $k < 0$ and $X(k)$ is finite-dimensional for all $k \geq 0$.

Let $(I, \leq)$ be a totally ordered index set, and for each $i \in I$ let X_i be a connected braided Hopf algebra in ${}^H_H\mathcal{YD}$. Let 1 denote the unit of (each) X_i . For any finite subset $J = \{j_1 < j_2 < \dots < j_t\} \subset I$ let $X_J = X_{j_1} \otimes X_{j_2} \otimes \dots \otimes X_{j_t}$. The family of objects X_J forms a direct system with respect to the inclusion in the following sense. If $J \subset K \subset I$ are finite subsets, where $J = \{j_1 < j_2 < \dots < j_t\}$, $K = \{k_1 < k_2 < \dots < k_s\}$, then $X_J \subset X_K$ via the embedding

$$X_J \rightarrow X_K, \quad x_{j_1} \otimes x_{j_2} \otimes \dots \otimes x_{j_t} \mapsto y_{k_1} \otimes y_{k_2} \otimes \dots \otimes y_{k_s},$$

where for all l with $1 \leq l \leq s$ let

$$y_{k_l} = \begin{cases} x_{j_m} & \text{if } k_l = j_m \text{ for some } m \leq t, \\ 1 & \text{otherwise.} \end{cases}$$

We write $\bigotimes_{i \in I} X_i$ for the limit of this direct system.

Remark 4.2. If X_i is $\mathbb{Z}^\theta$ -graded for all $i \in I$, then $\bigotimes_{i \in I} X_i$ is $\mathbb{Z}^\theta$ -graded. If $(X_i)_0 = \mathbb{k}$ and $X_i = \bigoplus_{\gamma \in \mathbb{N}_0\mathcal{A}} (X_i)_\gamma$ for all $i \in I$, then $\bigotimes_{i \in I} X_i \in {}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$ if and only if $X_i \in {}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$ for all $i \in I$ and for all $\gamma \in \mathbb{N}_0\mathcal{A} \setminus \{0\}$ the set $\{i \in I \mid (X_i)_\gamma \neq 0\}$ is finite.

Theorem 4.3. *Let R be a connected braided Hopf algebra in ${}^H_H\mathcal{YD}$, $\kappa \in \mathbb{N}$, and let $V_1, \dots, V_\kappa \in {}^H_H\mathcal{YD}$ be subobjects of $P(R)$, such that R is generated by $\bigoplus_{i=1}^\kappa V_i$. Let $\{e_1, \dots, e_\kappa\}$ be the standard basis of $\mathbb{Z}^\kappa$, and assume that there is an $\mathbb{N}_0^\kappa$ -grading of R such that $\deg V_i = e_i$ for all $i \in \{1, \dots, \kappa\}$. If for all $\mathbb{N}_0^\kappa$ -graded subalgebras $T \subset R$ and all $\mathbb{N}_0^\kappa$ -graded ideals T' of T there is an $\mathbb{N}_0^\kappa$ -graded splitting $T/T' \rightarrow R$ in ${}^H_H\mathcal{YD}$, then there exists a totally ordered index set $(L, \leq)$ and a family $(R_l)_{l \in L}$ of connected $\mathbb{N}_0^\kappa$ -graded braided Hopf algebras $R_l \in {}^H_H\mathcal{YD}$, such that*

- (1) $\{1, \dots, \kappa\} \subset L$, and for $l \in \{1, \dots, \kappa\}$, R_l is generated by an $\mathbb{N}_0^\kappa$ -homogeneous subspace $E_l \subset R_l$ of degree e_l such that $E_l \simeq V_l$ in ${}^H_H\mathcal{YD}$,
- (2) for all $l \in L \setminus \{1, \dots, \kappa\}$, R_l is generated by an $\mathbb{N}_0^\kappa$ -homogeneous subspace $E_l \in {}^H_H\mathcal{YD}$ of degree $\beta_l = \sum_{j=1}^\kappa n_{lj}e_j$, where at least two of the coefficients n_{lj} are nonzero,
- (3) $R \simeq \bigotimes_{l \in L} R_l$ as $\mathbb{N}_0^\kappa$ -graded objects in ${}^H_H\mathcal{YD}$.

Proof. See [GH07, Thm. 4.12]. The index set L corresponds to an appropriate subset of the set of Lyndon words in [GH07, Thm. 4.12]. By assumption, the maps ι_u in [GH07, Thm. 4.12] can be chosen to be $\mathbb{Z}^\kappa$ -homogeneous morphisms in ${}^H_H\mathcal{YD}$. Further, following the construction of R_l as subquotients of R and using the assumption that R is $\mathbb{N}_0^\kappa$ -graded, it is clear that all objects and morphisms in the theorem, including the direct limit $\bigotimes_{l \in L}$, are $\mathbb{N}_0^\kappa$ -graded. $\square$

The main technical tool in this section is the following lemma.

Lemma 4.4. *Let $0 \neq V \in {}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$ such that $V = \bigoplus_{\gamma \in \mathbb{N}_0 \setminus \{0\}} V_\gamma$, and all finite tensor powers of V are semisimple in ${}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$. Let $d_V = \min\{d \in \mathbb{N} \mid V(d) \neq 0\}$, $\kappa \in \mathbb{N}_{\geq 2}$ such that $V(d_V) = \bigoplus_{i=1}^{\kappa-1} V_i$ is a decomposition into irreducible objects in ${}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$, and let $V_\kappa = \bigoplus_{n=d_V+1}^\infty V(n)$. Then there is a totally ordered index set $(L, \leq)$ and a family $(W_l)_{l \in L}$, $0 \neq W_l \in {}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$ for all $l \in L$, such that*

- (1) $\{1, \dots, \kappa-1\} \subset L$ and $W_l \simeq V_l$ in ${}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$ for $l \in \{1, \dots, \kappa-1\}$,
- (2) if $l \in L \setminus \{1, 2, \dots, \kappa-1\}$, then $W_l = \bigoplus_{n=d_V+1}^\infty W_l(n) \in {}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$,
- (3) $\mathfrak{B}(V) \simeq \bigotimes_{l \in L} \mathfrak{B}(W_l)$ as objects in ${}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$.

Proof. Let H' be as in Rem. 4.1. We will apply Thm. 4.3 to objects in ${}^{H'}_H\mathcal{YD}$ instead of ${}^H_H\mathcal{YD}$.

Recall that $\mathfrak{B}(V)$ is a connected braided Hopf algebra in ${}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$ generated by $V \subset P(\mathfrak{B}(V))$. Let $\{e_i \mid 1 \leq i \leq \kappa\}$ be the standard basis of $\mathbb{Z}^\kappa$. The assignment

$$\underline{\deg} v = e_i, \quad \text{where } v \in V_i,$$

defines an $\mathbb{N}_0^\kappa$ -grading on V , and this extends to an $\mathbb{N}_0^\kappa$ -grading of $\mathfrak{B}(V)$ which is compatible with the $\mathbb{Z}^\theta$ -grading: $\mathbb{Z}^\theta$ -homogeneous components of a $\mathbb{Z}^\kappa$ -homogeneous element are $\mathbb{Z}^\kappa$ -homogeneous and *vice versa*. Thus we may regard $\mathfrak{B}(V)$ as a $\mathbb{Z}^\kappa$ -graded object in ${}^{H'}_H\mathcal{YD}$. Apply Thm. 4.3 to $\mathfrak{B}(V) \in {}^{H'}_H\mathcal{YD}$. This is possible, since the existence of the splittings $T/T' \rightarrow R$ in ${}^{H'}_H\mathcal{YD}$ follows from the assumption that finite tensor powers of V are semisimple in ${}^{H'}_H\mathcal{YD}$. We conclude that there exists a totally ordered index set $(L, \leq)$ and a family $(R_l)_{l \in L}$ of connected $\mathbb{Z}^{\theta+\kappa}$ -graded braided Hopf algebras $R_l \in {}^H_H\mathcal{YD}$, such that Thm. 4.3(1)–(3) hold. Since E_l is $\mathbb{N}_0^\kappa$ -homogeneous for all $l \in L$, the degrees of the $\mathbb{N}_0^\kappa$ -homogeneous components of R_l are of the form $n \underline{\deg} E_l$, where $n \in \mathbb{N}_0$. Further, Thm. 3.5 implies (use Rem. 4.1 with $\kappa + \theta$ instead of κ) that for all $l \in L$ there exists an $\mathbb{N}_0^\kappa$ -graded object $W_l \in {}^{H'}_H\mathcal{YD}$ such that $E_l \subset W_l$ and $R_l \simeq \mathfrak{B}(W_l)$ in ${}^{H'}_H\mathcal{YD}$. By the above,

- (*) the degrees of the $\mathbb{N}_0^\kappa$ -homogeneous components of W_l are multiples of $\underline{\deg} E_l$.

Then Thm. 4.3(3) and relation $\mathfrak{B}(V) \in {}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$ imply that

$$(**) \mathfrak{B}(V) \simeq \otimes_{l \in L} \mathfrak{B}(W_l) \text{ in } {}^H_H\mathcal{YD}^{\mathbb{Z}^{\theta+\kappa}} \text{ and in } {}^H_H\mathcal{YD}^{\mathbb{Z}^\theta},$$

and hence Claim (3) of the lemma holds.

Let first $l \in \{1, 2, \dots, \kappa - 1\}$. Then

$$\mathfrak{B}(V) = \mathfrak{B}(V_l \oplus \tilde{V}_l) = \mathfrak{B}(V_l) \oplus (\tilde{V}_l),$$

where $\tilde{V}_l = \oplus_{1 \leq j \leq \kappa, j \neq l} V_j$ and $(\tilde{V}_l) \subset \mathfrak{B}(V)$ is the ideal generated by $\tilde{V}_l$. Thus

$$\mathfrak{B}(V_l) = \oplus_{n \in \mathbb{N}_0} \{v \in \mathfrak{B}(V) \mid \underline{\deg} v = n e_l\}.$$

Further, $\mathfrak{B}(W_l)$ is a connected braided Hopf algebra generated by W_l , and $V_l \simeq E_l \subset W_l$. By (*), (**), and since the $\mathbb{N}_0^\kappa$ -homogeneous

components of $\mathfrak{B}(V_l)$ are finite-dimensional, it follows that $V_l \simeq W_l$ in ${}^H_H\mathcal{YD}^{\mathbb{Z}^{\theta+\kappa}}$. This gives (1).

Let now $l \in L \setminus \{1, 2, \dots, \kappa - 1\}$ and $W_{l\beta}$ an $\mathbb{N}_0^\kappa$ -homogeneous component of W_l . Then either $\beta = e_\kappa$ or $\beta = \sum_{j=1}^\kappa n_j e_j$ with $\sum_{j=1}^\kappa n_j \geq 2$. In the first case (**) implies that $W_{l\beta} \subset V_\kappa$, and hence $d_{W_l} \geq d_V + 1$. In the second case $d_{W_l} \geq 2d_V \geq d_V + 1$, since $d_V \geq 1$. Hence Claim (2) is proven. $\square$

Theorem 4.5. *Let $\theta \in \mathbb{N}$ and let $V_1, \dots, V_\theta$ be finite-dimensional irreducible objects in ${}^H_H\mathcal{YD}$. Let $V = \bigoplus_{i=1}^\theta V_i$, and assume that finite tensor powers of V are semisimple in ${}^H_H\mathcal{YD}$. Define a $\mathbb{Z}^\theta$ -grading of $\mathfrak{B}(V)$ such that $\deg V_i = \alpha_i$ for all i .*

(1) *There exists a totally ordered index set $(L, \leq)$ and a family $(W_l)_{l \in L}$ of irreducible objects $W_l \in {}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$ with $\deg W_l \in \mathbb{N}_0 \mathcal{A}$ for all l , and*

$$\mathfrak{B}(V) \simeq \bigotimes_{l \in L} \mathfrak{B}(W_l) \quad \text{in } {}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}.$$

(2) *If $\mathfrak{B}(V) \simeq \bigotimes_{l \in L} \mathfrak{B}(W_l)$ and $\mathfrak{B}(V) \simeq \bigotimes_{l' \in L'} \mathfrak{B}(W_{l'})$ for index sets $(L, \leq)$, $(L', \leq)$, and families $(W_l)_{l \in L}$, $(W_{l'})_{l' \in L'}$ as in (1), then there exists a bijection $\varphi : L \rightarrow L'$ such that $W_l \simeq W'_{\varphi(l)}$ in ${}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$ for all $l \in L$.*

Proof. The uniqueness follows from Lemma 4.7 below.

For the proof of the existence of the family $(W_l)_{l \in L}$ we first construct an inverse system of totally ordered sets and corresponding families of objects in ${}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$. This inverse system gives rise to a direct system in a natural way, and the limit of this direct system will be the family $(W_l)_{l \in L}$ we are looking for.

First we define recursively for all $k \in \mathbb{N}_0$ a totally ordered index set $(L^k, \leq)$ and a family $(W_l^k)_{l \in L^k}$, such that

$$(*) \quad 0 \neq W_l^k \in {}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}, \quad W_l^k \text{ is irreducible or } W_l^k = \bigoplus_{n=k+1}^\infty W_l^k(n) \\ \text{for all } l \in L^k, \text{ and } \mathfrak{B}(V) \simeq \bigotimes_{l \in L^k} \mathfrak{B}(W_l^k) \text{ in } {}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}.$$

Note that then for all k, l, n , $W_l^k(n)$ is isomorphic to a direct summand of $V^{\otimes n}$, and hence all finite tensor powers of W_l^k are semisimple in ${}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$.

For $k = 0$ let $L^0 = \{1\}$ and $W_1^0 = V$. If $k \geq 0$, $l \in L^k$, and W_l^k is not irreducible, then we choose a totally ordered index set $(L_{kl}, \leq)$ and a family $(W_{klm})_{m \in L_{kl}}$ of objects in ${}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$ such that

$$\mathfrak{B}(W_l^k) \simeq \bigotimes_{m \in L_{kl}} \mathfrak{B}(W_{klm})$$

and $0 \neq W_{klm} \in {}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$ is irreducible or $W_{klm} = \bigoplus_{n=k+2}^\infty W_{klm}(n)$. This is possible by Lemma 4.4. Now we define

$$(4.1) \quad \begin{aligned} L'^k &= \{l \in L^k \mid W_l^k \text{ is irreducible}\}, \\ L^{k+1} &= L'^k \sqcup \{(l, m) \mid l \in L^k \setminus L'^k, m \in L_{kl}\}, \quad (\text{disjoint union}) \\ W_j^{k+1} &= \begin{cases} W_j^k & \text{if } j \in L'^k, \\ W_{klm} & \text{if } j = (l, m), l \in L^k \setminus L'^k, m \in L_{kl}. \end{cases} \end{aligned}$$

Let $\leq$ be the natural order on L^{k+1} . That is, if $l_1, l_2 \in L'^k$, then $l_1 \leq l_2$ if and only if this relation holds in L^k . If $l_1 \in L'^k$, $l_2 = (\tilde{l}, m) \in L^{k+1} \setminus L'^k$, then $l_1 \leq l_2$ if and only if $l_1 \leq \tilde{l}$ in L^k , and $l_2 \leq l_1$ if and only if $\tilde{l} \leq l_1$ in L^k . Finally, if $l_1 = (\tilde{l}_1, m_1)$, $l_2 = (\tilde{l}_2, m_2) \in L^{k+1} \setminus L'^k$, then $l_1 \leq l_2$ if and only if $\tilde{l}_1 < \tilde{l}_2$ in L^k or $\tilde{l}_1 = \tilde{l}_2$ and $m_1 \leq m_2$ in $L_{k\tilde{l}_1}$. Clearly, $(L^{k+1}, \leq)$ is totally ordered, and the family $(W_l^{k+1})_{l \in L^{k+1}}$ satisfies the properties in (*).

For all $k \in \mathbb{N}_0$ define L'^k as in Eq. (4.1), and let $L = \bigcup_{k \in \mathbb{N}} L'^k$ with the total order $\leq$ induced by the orders defined on the sets L'^k . Note that $L^k \subset L'^{k+1}$ and $W_l^{k+1} = W_l^k$ for all $k \in \mathbb{N}_0$ and $l \in L'^k$. Define $W_l = W_l^k$ for all $l \in L'^k \subset L$. Then $0 \neq W_l \in {}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$ is irreducible and has positive degree (with respect to the basis $\{\alpha_i \mid 1 \leq i \leq \theta\}$) for all $l \in L$. We prove that $\bigotimes_{l \in L} \mathfrak{B}(W_l) \simeq \mathfrak{B}(V)$ in ${}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$. Since $L'^k \subset L^k$ for all $k \in \mathbb{N}_0$, $\bigotimes_{l \in L'^k} \mathfrak{B}(W_l)$ is isomorphic to a subobject of $\mathfrak{B}(V)$ for all $k \in \mathbb{N}_0$ by (*). Moreover, (*) also implies that

$$(4.2) \quad \mathfrak{B}^n(V) \simeq \left(\bigotimes_{l \in L'^k} \mathfrak{B}(W_l) \right)(n)$$

for all $k, n \in \mathbb{N}_0$ with $n \leq k$. The construction of L and $(W_l)_{l \in L}$ shows that

$$\left(\bigotimes_{l \in L'^k} \mathfrak{B}(W_l) \right)(n) = \left(\bigotimes_{l \in L'^{k+1}} \mathfrak{B}(W_l) \right)(n) = \left(\bigotimes_{l \in L} \mathfrak{B}(W_l) \right)(n)$$

for all $k, n \in \mathbb{N}_0$ and $n \leq k$. Hence

$$\begin{aligned} \mathfrak{B}(V) &= \oplus_{n=0}^{\infty} \mathfrak{B}^n(V) \simeq \oplus_{k=0}^{\infty} (\otimes_{l \in L'^k} \mathfrak{B}(W_l))(k) \\ &= \oplus_{k=0}^{\infty} (\otimes_{l \in L} \mathfrak{B}(W_l))(k) = \otimes_{l \in L} \mathfrak{B}(W_l). \end{aligned}$$

This finishes the proof of the theorem. $\square$

Remark 4.6. In the setting of Thm. 4.5, assume that H is the group algebra of an abelian group, and V_i is one-dimensional for all $i \in \{1, 2, \dots, \theta\}$. In [Kha99], Kharchenko gave a construction of a PBW-basis for a class of Hopf algebras, which can be applied to $\mathfrak{B}(V)$. Thm. 4.5 can be viewed as a generalization of Kharchenko's basis to Nichols algebras over arbitrary Hopf algebras in the sense that the PBW generators of Kharchenko are replaced by the Yetter-Drinfeld modules W_l .

Lemma 4.7. *Let $\theta \in \mathbb{N}$, $(L, \leq)$, $(L', \leq)$ totally ordered index sets, and $(W_l)_{l \in L}$, $(W'_{l'})_{l' \in L'}$ families of irreducible objects in ${}^H_H \mathcal{YD}^{\mathbb{Z}^\theta}$. Assume that $\deg W_l \in \sum_{i=1}^{\theta} \mathbb{N}_0 \alpha_i \setminus \{0\}$ for all $l \in L$. If $\otimes_{l \in L} \mathfrak{B}(W_l) \simeq \otimes_{l' \in L'} \mathfrak{B}(W'_{l'})$, and all $\mathbb{Z}^\theta$ -homogeneous components of $\otimes_{l \in L} \mathfrak{B}(W_l)$ are finite-dimensional, then there exists a bijection $\varphi : L \rightarrow L'$ such that $W_l \simeq W'_{\varphi(l)}$ in ${}^H_H \mathcal{YD}^{\mathbb{Z}^\theta}$ for all $l \in L$.*

Proof. Let $l' \in L'$. Since $W'_{l'}$ is a direct summand of $\otimes_{l'' \in L'} \mathfrak{B}(W'_{l''}) \simeq \otimes_{l \in L} \mathfrak{B}(W_l)$ and the degree 0 component of $\otimes_{l \in L} \mathfrak{B}(W_l)$ is $\mathbb{k}$, we obtain that $\deg W'_{l'} \in \sum_{i=1}^{\theta} \mathbb{N}_0 \alpha_i \setminus \{0\}$.

For $k \in \mathbb{N}_0$ let $L_k = \{l \in L \mid |\deg W_l| = k\}$. Then $L_0 = \emptyset$, $L = \cup_{k \in \mathbb{N}} L_k$, and L_k is a finite set for all $k \in \mathbb{N}$, since the $\mathbb{Z}^\theta$ -homogeneous components of $\otimes_{l \in L} \mathfrak{B}(W_l)$ are finite-dimensional. Similarly $L' = \cup_{k \in \mathbb{N}} L'_k$. It suffices to show that

- (*) for all $k \in \mathbb{N}_0$ there is a bijection $\varphi_k : L_k \rightarrow L'_k$ such that $W_l \simeq W'_{\varphi_k(l)}$ in ${}^H_H \mathcal{YD}^{\mathbb{Z}^\theta}$ for all $k \in \mathbb{N}_0$, $l \in L_k$.

This is clear for $k = 0$, since $L_0 = L'_0 = \emptyset$.

Let $k \in \mathbb{N}$. Then

$$\begin{aligned} (\otimes_{l \in L} \mathfrak{B}(W_l))(k) &\simeq \oplus_{l \in L_k} W_l \oplus (\otimes_{l \in L_1 \cup L_2 \cup \dots \cup L_{k-1}} \mathfrak{B}(W_l))(k), \\ (\otimes_{l' \in L'} \mathfrak{B}(W'_{l'}))(k) &\simeq \oplus_{l' \in L'_k} W'_{l'} \oplus (\otimes_{l' \in L'_1 \cup L'_2 \cup \dots \cup L'_{k-1}} \mathfrak{B}(W'_{l'}))(k) \end{aligned}$$

in ${}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$. Hence $(*)$ follows by induction on k and by Krull-Remak-Schmidt. $\square$

The following similar results will be used later on.

Lemma 4.8. *Let K, K' and B be $\mathbb{Z}^\theta$ -graded objects in ${}^H_H\mathcal{YD}$ such that all homogeneous components of K and B are finite-dimensional, B_0 is isomorphic to the trivial object $\mathbb{k} \in {}^H_H\mathcal{YD}$, and $K_\gamma = B_\gamma = 0$ for all $\gamma \in \mathbb{Z}^\theta \setminus \sum_{i=1}^\theta \mathbb{N}_0 \alpha_i$. If $K \otimes B \simeq K' \otimes B$ in ${}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$, then $K \simeq K'$ in ${}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$.*

Proof. Since $B_0 \simeq \mathbb{k}$, K'_γ is (isomorphic to) a direct summand of $(K' \otimes B)_\gamma$ for all $\gamma \in \mathbb{Z}^\theta$. Hence the isomorphism $K \otimes B \simeq K' \otimes B$ implies that $K'_\gamma = 0$ for all $\gamma \in \mathbb{Z}^\theta \setminus \sum_{i=1}^\theta \mathbb{N}_0 \alpha_i$.

Similarly to the proof of the previous lemma,

$$\begin{aligned} (K \otimes B)(k) &\simeq K(k) \oplus \bigoplus_{1 \leq k' \leq k} K(k - k') \otimes B(k'), \\ (K' \otimes B)(k) &\simeq K'(k) \oplus \bigoplus_{1 \leq k' \leq k} K'(k - k') \otimes B(k') \end{aligned}$$

in ${}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$ for all $k \in \mathbb{N}_0$. Since the $\mathbb{N}_0$ -homogeneous components of $K \otimes B$ are finite-dimensional, the claim follows by induction on k and by Krull-Remak-Schmidt. $\square$

Lemma 4.9. *Let $\theta \in \mathbb{N}$ and let $V_1, \dots, V_\theta$ be finite-dimensional irreducible objects in ${}^H_H\mathcal{YD}$. Let $V = \bigoplus_{i=1}^\theta V_i$, and define a $\mathbb{Z}^\theta$ -grading of $\mathfrak{B}(V)$ such that $\deg V_i = \alpha_i$ for all i . Assume that there exists a totally ordered index set $(L, \leq)$ and a family $(W_l)_{l \in L}$ of irreducible objects $W_l \in {}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$ such that $\deg W_l \in \mathbb{N}_0 \mathcal{A}$ for all l , and*

$$\mathfrak{B}(V) \simeq \bigotimes_{l \in L} \mathfrak{B}(W_l)$$

in ${}^H_H\mathcal{YD}^{\mathbb{Z}^\theta}$. Then for each $i \in \{1, \dots, \theta\}$ there exists a unique $l(i) \in L$ such that $\deg W_{l(i)} \in \mathbb{N} \alpha_i$. Moreover, $\deg W_{l(i)} = \alpha_i$ and $W_{l(i)} \simeq V_i$.

Proof. Let $i \in \{1, \dots, \theta\}$. The subspace $\bigoplus_{n \in \mathbb{Z}} \mathfrak{B}(V)_{n\alpha_i} \subset \mathfrak{B}(V)$ is the subalgebra $\mathfrak{B}(V_i)$, and $\mathfrak{B}(V)_{\alpha_i} = V_i$. Since $\deg W_l \in \sum_{j=1}^\theta \mathbb{N}_0 \alpha_j \setminus 0$, we obtain that

$$(4.3) \quad \mathfrak{B}(V_i) \simeq \bigoplus_{n \in \mathbb{Z}} \left(\bigotimes_{l \in L} \mathfrak{B}(W_l) \right)_{n\alpha_i} = \bigotimes_{l \in L_i} \mathfrak{B}(W_l),$$

where $L_i = \{l \in L \mid \deg W_l \in \mathbb{N}\alpha_i\}$. Hence

$$V_i = \mathfrak{B}(V_i)_{\alpha_i} \simeq \left(\bigotimes_{l \in L} \mathfrak{B}(W_l) \right)_{\alpha_i} = \bigoplus_{l \in L_i, \deg W_l = \alpha_i} W_l,$$

and since V_i is irreducible, it follows that there is a unique $l = l(i) \in L$ such that $V_i \simeq W_l$ and $\deg W_l = \alpha_i$. Then Eq. (4.3) implies that $L_i = \{l(i)\}$ which proves the lemma. $\square$

5. WEYL GROUPOIDS AND ROOT SYSTEMS

In this section we give the definition of generalized root systems in [HY08] in terms of category theory, see [CH08], and recall some results which will be needed in the following sections.

Let I be a non-empty finite set and $(\alpha_i)_{i \in I}$ the standard basis of $\mathbb{Z}^I$.

Recall from [Kac90, §1.1] that a generalized Cartan matrix $A = (a_{ij})_{i,j \in I}$ is a matrix in $\mathbb{Z}^{I \times I}$ such that

- (M1) $a_{ii} = 2$ and $a_{jk} \leq 0$ for all $i, j, k \in I$ with $j \neq k$,
- (M2) if $i, j \in I$ and $a_{ij} = 0$, then $a_{ji} = 0$.

Let $\mathcal{M}$ be a non-empty set, and for all $i \in I$ and $N \in \mathcal{M}$ let $r_i : \mathcal{M} \rightarrow \mathcal{M}$ be a map and $A^N = (a_{jk}^N)_{j,k \in I}$ a generalized Cartan matrix. The quadruple

$$\mathcal{C} = \mathcal{C}(I, \mathcal{M}, (r_i)_{i \in I}, (A^N)_{N \in \mathcal{M}}),$$

is called a *Cartan scheme* if

- (C1) $r_i^2 = \text{id}$ for all $i \in I$,
- (C2) $a_{ij}^N = a_{ij}^{r_i(N)}$ for all $N \in \mathcal{M}$ and $i, j \in I$.

Let $\mathcal{C} = \mathcal{C}(I, \mathcal{M}, (r_i)_{i \in I}, (A^N)_{N \in \mathcal{M}})$ be a Cartan scheme. For all $i \in I$, $N \in \mathcal{M}$ define $s_i^N \in \text{Aut}(\mathbb{Z}^I)$ by

$$s_i^N(\alpha_j) = \alpha_j - a_{ij}^N \alpha_i \quad \text{for all } j \in I.$$

Recall that a groupoid is a category where all morphisms are isomorphisms. The *Weyl groupoid of $\mathcal{C}$* is the groupoid $\mathcal{W}(\mathcal{C})$ with $\text{Ob}(\mathcal{W}(\mathcal{C})) = \mathcal{M}$, where the morphisms are generated by all $s_i^N : N \rightarrow r_i(N)$ with $i \in I$, $N \in \mathcal{M}$. Formally, for $N, P \in \mathcal{M}$ the set $\text{Hom}(N, P)$ consists of the triples (P, s, N) , such that

$$s = s_{i_n}^{r_{i_{n-1}} \cdots r_{i_1}(N)} \cdots s_{i_2}^{r_{i_1}(N)} s_{i_1}^N$$

and $r_{i_n} \cdots r_{i_2} r_{i_1}(N) = P$ for some $n \in \mathbb{N}_0$ and $i_1, \dots, i_n \in I$. The composition of morphisms is induced by the group structure of $\text{Aut}(\mathbb{Z}^I)$:

$$(Q, g, P) \circ (P, f, N) = (Q, gf, N)$$

for all $(Q, g, P), (P, f, N) \in \text{Hom}(\mathcal{W}(\mathcal{C}))$. If $w = (P, f, N) \in \text{Hom}(\mathcal{W}(\mathcal{C}))$ and $\alpha \in \mathbb{Z}^\theta$, then we define $w(\alpha) = f(\alpha)$.

Note that the inverse of the generating morphism $(r_i(N), s_i^N, N)$ is $(N, s_i^{r_i(N)}, r_i(N))$, since s_i^N is a reflection, and $s_i^N = s_i^{r_i(N)}$ and $r_i^2(N) = N$ by definition.

We say that

$$\mathcal{R} = \mathcal{R}(\mathcal{C}, (\Delta(N))_{N \in \mathcal{M}})$$

is a *root system of type $\mathcal{C}$* if $\mathcal{C} = \mathcal{C}(I, \mathcal{M}, (r_i)_{i \in I}, (A^N)_{N \in \mathcal{M}})$ is a Cartan scheme and $\Delta(N) \subset \mathbb{Z}^I$, where $N \in \mathcal{M}$, are subsets such that

- (R1) $\Delta(N) = (\Delta(N) \cap \mathbb{N}_0^I) \cup -(\Delta(N) \cap \mathbb{N}_0^I)$ for all $N \in \mathcal{M}$,
- (R2) $\Delta(N) \cap \mathbb{Z}\alpha_i = \{\alpha_i, -\alpha_i\}$ for all $i \in I, N \in \mathcal{M}$,
- (R3) $s_i^N(\Delta(N)) = \Delta(r_i(N))$ for all $i \in I, N \in \mathcal{M}$,
- (R4) $(r_i r_j)^{m_{i,j;N}}(N) = N$ for all $i, j \in I$ and $N \in \mathcal{M}$ such that $i \neq j$ and $m_{i,j;N} := |\Delta(N) \cap (\mathbb{N}_0\alpha_i + \mathbb{N}_0\alpha_j)|$ is finite.

We note that Axiom (M2) is redundant for root systems by [CH08, Lemma 2.5].

If $\mathcal{R}(\mathcal{C}, (\Delta(N))_{N \in \mathcal{M}})$ is a root system of type $\mathcal{C}$, then $\mathcal{W}(\mathcal{R}) := \mathcal{W}(\mathcal{C})$ is called the *Weyl groupoid of $\mathcal{R}$* . The elements of $\Delta_+(N) := \Delta(N) \cap \mathbb{N}_0^I$ and $\Delta_-(N) := -\Delta_+(N)$ are called *positive* and *negative roots*, respectively. Following [Kac90, §5.1] we say that the roots $w(\alpha_i) \in \Delta(P)$, where $w \in \text{Hom}(N, P)$, $N, P \in \mathcal{M}$, and $i \in I$, are *real roots*. The set of real roots and positive real roots is denoted by $\Delta^{\text{re}}(N)$ and $\Delta_+^{\text{re}}(N)$, respectively. Note that (R3) implies that $w(\Delta(N)) = \Delta(P)$ for all $N, P \in \text{Ob}(\mathcal{W}(\mathcal{R}))$ and $w \in \text{Hom}(N, P)$.

Recall that a groupoid G is *connected*, if for all $N, P \in \text{Ob}(G)$ the set $\text{Hom}(N, P)$ is non-empty. It is *finite*, if $\text{Hom}(G)$ is finite.

Lemma 5.1. [CH08, Lemma 2.11] *Let $\mathcal{C}$ be a Cartan scheme and $\mathcal{R}$ a root system of type $\mathcal{C}$. Assume that $\mathcal{W}(\mathcal{R})$ is connected. Then the following are equivalent.*

- (1) $\Delta(N)$ is finite for all $N \in \text{Ob}(\mathcal{W}(\mathcal{R}))$.

- (2) $\Delta(N)$ is finite for at least one $N \in \text{Ob}(\mathcal{W}(\mathcal{R}))$.
- (3) $\Delta^{\text{re}}(N)$ is finite for all $N \in \text{Ob}(\mathcal{W}(\mathcal{R}))$.
- (4) $\mathcal{W}(\mathcal{R})$ is finite.

Proposition 5.2. [CH08, Prop. 2.12] *Let $\mathcal{C}$ be a Cartan scheme and $\mathcal{R}$ a root system of type $\mathcal{C}$. If $\mathcal{W}(\mathcal{R})$ is finite, then all roots are real.*

Lemma 5.3. [HY08, Lemma 8(iii)] *Let $\mathcal{C}$ be a Cartan scheme and $\mathcal{R}$ a root system of type $\mathcal{C}$. Let $N, P \in \text{Ob}(\mathcal{W}(\mathcal{R}))$ and $w \in \text{Hom}(N, P) \subset \text{Hom}(\mathcal{W}(\mathcal{R}))$ such that $w(\Delta_+(N)) \subset \Delta_-(P)$. Then $\Delta(N)$ is finite.*

In general the Cartan matrices A^N are not of finite type if the Weyl groupoid $\mathcal{W}(\mathcal{R})$ is finite [CH08, Prop. 5.1(2)]. But in the special case, when the Cartan scheme $\mathcal{C}$ is *standard*, i.e. $A^N = A^M$ for all $N, M \in \text{Ob}(\mathcal{W}(\mathcal{C}))$, the following holds (see [CH08, Thm. 3.3] for a slightly different proof).

Corollary 5.4. *Let $\mathcal{C}$ be a standard Cartan scheme with generalized Cartan matrix $A = A^N$ for all $N \in \text{Ob}(\mathcal{W}(\mathcal{C}))$. Let $\mathcal{R}$ be a root system of type $\mathcal{C}$. Then the following are equivalent.*

- (1) $\mathcal{W}(\mathcal{R})$ is finite.
- (2) A is a Cartan matrix of finite type.

Proof. Let $W(A)$ be the Weyl group of the Kac-Moody Lie algebra of A . In particular, $W(A)$ is isomorphic to the subgroup of $\text{Aut}(\mathbb{Z}^\theta)$ generated by the reflections $s_i \in \text{Aut}(\mathbb{Z}^\theta)$, where

$$s_i(\alpha_j) = \alpha_j - a_{ij}\alpha_i \quad \text{for all } i, j \in I.$$

(1) $\Rightarrow$ (2): Since $\mathcal{C}$ is standard, the map $\text{Hom}(\mathcal{W}(\mathcal{R})) \rightarrow W(A)$, $(P, s, N) \mapsto s$, is well-defined and surjective. Hence $W(A)$ is finite by (1), and A is of finite type.

(2) $\Rightarrow$ (1): Since A is of finite type, the Weyl group $W(A)$ acts transitively on bases of the root system. Hence there exists a permutation τ of I and an element $w \in W(A)$ such that $w(\alpha_i) = -\alpha_{\tau(i)}$ for all $i \in I$. Let $n \in \mathbb{N}$ and $i_1, \dots, i_n \in I$ such that $w = s_{i_n} \cdots s_{i_2} s_{i_1}$. Then $w = s_{i_n}^{r_{i_n-1} \cdots r_{i_1}(N)} \cdots s_{i_2}^{r_{i_1}(N)} s_{i_1}^N$ for all $N \in \mathcal{M}$, since $\mathcal{C}$ is standard. Hence $w(\Delta_+(N)) \subset \Delta_-(r_{i_n} \cdots r_{i_2} r_{i_1}(N))$ for all $N \in \mathcal{M}$. This proves (1) by Lemmata 5.1 and 5.3. $\square$

Remark 5.5. Let $\mathcal{C} = \mathcal{C}(I, \mathcal{M}, (r_i)_{i \in I}, (A^N)_{N \in \mathcal{M}})$ be a standard Cartan scheme and $\mathcal{R} = \mathcal{R}(\mathcal{C}, (\Delta^N)_{N \in \mathcal{M}})$ a root system of type $\mathcal{C}$. Define $A = A^N$ for all $N \in \mathcal{M}$, and let $W(A)$ denote the Weyl group of the Kac-Moody Lie algebra of A . Then the maps $s_i^N \in \text{Aut}(\mathbb{Z}^\theta)$, where $i \in I$ and $N \in \mathcal{M}$, are independent of N . Assume that $\Delta(N) = \Delta^{\text{re}}(N)$ for all $N \in \mathcal{M}$. Then $\Delta(N) = \{w(\alpha_i) \mid w \in W(A), i \in I\}$ for all $N \in \mathcal{M}$, and hence it is independent of N . In this case $\mathcal{R}$ is equivalent to an action of $W(A)$ on $\mathcal{M}$.

There is an important necessary condition for the finiteness of the Weyl groupoid of a Cartan scheme. For the proof we need a lemma, which is a special case of [Hec08, Lemma 9].

Lemma 5.6. *Let $T \subset \text{SL}(2, \mathbb{Z})$ be a non-empty subsemigroup generated by matrices of the form $\begin{pmatrix} a & -b \\ c & -d \end{pmatrix}$ with $0 < d < b < a$. Then all elements of T are of this form. In particular, $\text{id} \notin T$.*

Proposition 5.7. *Let $\mathcal{C} = \mathcal{C}(I, \mathcal{M}, (r_i)_{i \in I}, (A^N)_{N \in \mathcal{M}})$ be a Cartan scheme of rank two. If $\text{Hom}(\mathcal{W}(\mathcal{C}))$ is a finite set, then there exist $N \in \mathcal{M}$ and $i, j \in I$ such that $a_{ij}^N \in \{0, -1\}$.*

Proof. For all $a \in \mathbb{Z}$ let

$$\eta_1(a) = \begin{pmatrix} -1 & a \\ 0 & 1 \end{pmatrix}, \quad \eta_2(a) = \begin{pmatrix} 1 & 0 \\ a & -1 \end{pmatrix}.$$

If $N \in \mathcal{M}_2(M)$, then

$$(5.1) \quad \begin{aligned} & s_1^{r_2 \cdots r_1 r_2(M)} \cdots s_2^{r_1 r_2(M)} s_1^{r_2(M)} s_2^M \\ &= \eta_1(-a_{12}^{r_2 \cdots r_1 r_2(M)}) \cdots \eta_2(-a_{21}^{r_1 r_2(M)}) \eta_1(-a_{12}^{r_2(M)}) \eta_2(-a_{21}^M). \end{aligned}$$

Assume that $a_{12}^N, a_{21}^N < -1$ for all $N \in \mathcal{M}_2(M)$. Then the matrices

$$\eta_1(a') \eta_2(a'') = \begin{pmatrix} a'a'' - 1 & -a' \\ a'' & -1 \end{pmatrix},$$

where $a' = -a_{12}^{r_2(N)}$, $a'' = -a_{21}^N$ for some $N \in \mathcal{M}_2(M)$, satisfy the assumption of Lemma 5.6. Lemma 5.6 implies that the set of products given in Eq. (5.1) is infinite. This is a contradiction to the finiteness of $\text{Hom}(\mathcal{W}(\mathcal{C}))$. $\square$

6. WEYL GROUPOIDS AND ROOT SYSTEMS FOR NICHOLS ALGEBRAS

Let $\widehat{{}^H_H\mathcal{YD}}$ denote the set of isomorphism classes of finite-dimensional irreducible Yetter-Drinfeld modules over H . For any $X \in {}^H_H\mathcal{YD}$ let $[X]$ denote the isomorphism class of X . Let $\theta \in \mathbb{N}$, $\mathbb{I} = \{1, 2, \dots, \theta\}$, and $\mathcal{M}_\theta = (\widehat{{}^H_H\mathcal{YD}})^\theta$. The standard basis of $\mathbb{Z}^\theta$ is denoted by $\{\alpha_1, \dots, \alpha_\theta\}$.

Definition 6.1. Let $N \in \mathcal{M}_\theta$. Choose $N_1, \dots, N_\theta \in {}^H_H\mathcal{YD}$ such that $N = ([N_1], \dots, [N_\theta])$. Define $V(N) = N_1 \oplus \dots \oplus N_\theta$. Assume that there exists a totally ordered index set $(L, \leq)$ and a family $(W_l)_{l \in L}$ of finite-dimensional irreducible $\mathbb{N}_0^\theta$ -graded objects in ${}^H_H\mathcal{YD}$ such that $\mathfrak{B}(V(N)) \simeq \otimes_{l \in L} \mathfrak{B}(W_l)$ as $\mathbb{N}_0^\theta$ -graded objects in ${}^H_H\mathcal{YD}$, where $\deg N_i = \alpha_i$ for all $i \in \mathbb{I}$. Define

$$(6.1) \quad \begin{aligned} \tilde{\Delta}_+(N) = \{ (C, \alpha) \mid C \in \widehat{{}^H_H\mathcal{YD}}, \alpha \in \mathbb{N}_0^\theta \setminus 0, \\ \exists l \in L : C = [W_l], \alpha = \deg W_l \}, \end{aligned}$$

$$(6.2) \quad \tilde{\Delta}_-(N) = \{ (C^*, -\alpha) \mid (C, \alpha) \in \tilde{\Delta}_+(N) \},$$

$$(6.3) \quad \tilde{\Delta}(N) = \tilde{\Delta}_+(N) \cup \tilde{\Delta}_-(N),$$

and for all $(C, \alpha) \in \tilde{\Delta}_+(N)$ let

$$(6.4) \quad \text{mult}_N(C, \alpha) = \text{mult}_N(C^*, -\alpha) = |\{l \in L \mid C = [W_l], \alpha = \deg W_l\}|.$$

The sets

$$(6.5) \quad \Delta_+(N) = \{\deg W_l \mid l \in L\}, \quad \Delta_-(N) = -\Delta_+(N),$$

$$(6.6) \quad \Delta(N) = \Delta_+(N) \cup \Delta_-(N)$$

are called the *set of positive roots of N* , the *set of negative roots of N* , and the *set of roots of N* , respectively. For any $N \in \mathcal{M}_\theta$ we say that $\tilde{\Delta}(N)$ is defined, if the assumptions of Def. 6.1 are fulfilled for N .

Remark 6.2. In [Hec06b], the root system of a Nichols algebra of diagonal type was defined using Kharchenko's PBW-basis. Viewing the Yetter-Drinfeld modules W_l as generalized PBW generators, see Rem. 4.6, the definition of $\Delta(N)$ in Eq. (6.6) generalizes the root system given in [Hec06b].

Let $N \in \mathcal{M}_\theta$. By Thm. 4.5(1), $\tilde{\Delta}(N)$ is defined, if all tensor powers of $V(N)$ are semisimple.

By Lemma 4.7, the definitions of $\tilde{\Delta}_\pm(N)$, $\tilde{\Delta}(N)$, $\Delta_\pm(N)$, and $\Delta(N)$ only depend on N , but not on the choice of $N_1, \dots, N_\theta$, $(L, \leq)$, and the family $(W_l)_{l \in L}$.

Lemma 6.3. *Let $N, P \in \mathcal{M}_\theta$ such that $\tilde{\Delta}(N)$ and $\tilde{\Delta}(P)$ are defined. If $\tilde{\Delta}(N) = \tilde{\Delta}(P)$ then $N = P$.*

Proof. See Lemma 4.7. □

Definition 6.4. Let $N = ([N_1], \dots, [N_\theta]) \in \mathcal{M}_\theta$, $V(N) = N_1 \oplus \dots \oplus N_\theta$, and $i \in \mathbb{I}$. We say that N is *i-finite*, if for all $j \in \mathbb{I} \setminus \{i\}$ $(\text{ad}_c N_i)^h(N_j) = 0$ for some $h \in \mathbb{N}$, and *l-finite*, if N is *l-finite* for all $l \in \mathbb{I}$.

If N is not *i-finite*, then let $r_i(N) = N$.

Assume now that N is *i-finite*. Let $A^N = (a_{ij}^N)_{i,j \in \mathbb{I}}$, where

$$(6.7) \quad -a_{ij}^N = \sup\{h \in \mathbb{N}_0 \mid (\text{ad}_c N_i)^h(N_j) \neq 0 \text{ in } \mathfrak{B}(V(N))\}$$

for $j \in \mathbb{I} \setminus \{i\}$, and $a_{ii}^N = 2$. Let $r_i(N) = ([N'_1], \dots, [N'_\theta]) \in \mathcal{M}_\theta$, where $N'_i = N_i^*$ and $N'_j = (\text{ad}_c N_i)^{-a_{ij}^N}(N_j)$ for all $j \in \mathbb{I} \setminus \{i\}$. Define $s_i^N \in \text{Aut}(\mathbb{Z}^\theta)$ by

$$(6.8) \quad s_i^N(\alpha_j) = \alpha_j - a_{ij}^N \alpha_i \quad \text{for all } j \in \mathbb{I},$$

$$(6.9) \quad s_i^N(C, \gamma) = (C, s_i^N(\gamma)) \quad \text{for all } C \in \widehat{{}^H_H \mathcal{YD}}, \gamma \in \mathbb{Z}^\theta.$$

By definition, the map s_i^N is a reflection in the sense of [Bou68, Ch. V, §2.2], since $a_{ii}^N = 2$. Note that the definition of r_i is slightly different from the definition of $\mathcal{R}_i$ in [AHS08, Eq. (3.16)].

The next proposition shows how to compute the Yetter-Drinfeld module $(\text{ad}_c V)^n(W)$, where $n \in \mathbb{N}$ and $V, W \in {}^H_H \mathcal{YD}$. Recall the definition of S_n from Eq. (1.3).

Proposition 6.5. *Let $n \in \mathbb{N}$ and $V, W \in {}^H_H \mathcal{YD}$. Then the image of the linear map $(S_n \otimes \text{id})T_n \in \text{End}(V^{\otimes n} \otimes W)$, where*

$$T_n = (\text{id} - c_{n,n+1}^2 c_{n-1,n} \cdots c_{12}) \cdots (\text{id} - c_{n,n+1}^2 c_{n-1,n})(\text{id} - c_{n,n+1}^2),$$

is isomorphic to $(\text{ad}_c V)^n(W) \subset \mathfrak{B}(V \oplus W)$ in ${}^H_H \mathcal{YD}$.

Proof. The kernel of $S_{n+1} \in \text{End}((V \oplus W)^{\otimes n+1})$ is $\mathfrak{I}_{T(V \oplus W)}(n+1)$, see Sect. 1. Hence the image of S_{n+1} is isomorphic to $\mathfrak{B}(V \oplus W)(n+1)$ in ${}^H_H\mathcal{YD}$. The lemma follows from

$$(6.10) \quad (S_n \otimes \text{id})T_n(v_1 \otimes \cdots \otimes v_n \otimes w) = S_{n+1}((\text{ad}_c v_1) \cdots (\text{ad}_c v_n)(w))$$

for all $v_1, \dots, v_n \in V$, $w \in W$, where

$$\text{ad}_c v : (V \oplus W)^{\otimes k} \rightarrow (V \oplus W)^{\otimes k+1}, \quad x \mapsto v \otimes x - (v_{(-1)} \cdot x) \otimes v_{(0)}$$

for all $k \in \mathbb{N}_0$, $v \in V$, and $x \in (V \oplus W)^{\otimes k}$. By Eq. (1.3) it suffices to show that

$$T_n(v_1 \otimes \cdots \otimes v_n \otimes w) = S_{n,1}((\text{ad}_c v_1) \cdots (\text{ad}_c v_n)(w))$$

for all $v_1, \dots, v_n \in V$, $w \in W$. This follows by induction on n from the braid equation for c . $\square$

Let $N \in \mathcal{M}_\theta$ and $i \in \mathbb{I}$. Let $\mathcal{K}_i^N = \mathfrak{B}(V(N))^{\text{co } \mathfrak{B}(N_i)}$ be the algebra of right coinvariant elements with respect to the coaction $(\text{id} \otimes \pi) \Delta_{\mathfrak{B}(V(N))}$, where $\pi : \mathfrak{B}(V(N)) \rightarrow \mathfrak{B}(N_i)$ is the canonical projection, see [AHS08, Sect. 3.2]. Further, $\mathfrak{B}(V(N))$, $\mathcal{K}_i^N$, and $\mathfrak{B}(N_i)$ are $\mathbb{Z}^\theta$ -graded objects in ${}^H_H\mathcal{YD}$ with $\deg N_j = \alpha_j$ for all $j \in \mathbb{I}$. Then

$$(6.11) \quad \mathfrak{B}(V(N)) \simeq \mathcal{K}_i^N \otimes \mathfrak{B}(N_i)$$

as $\mathbb{N}_0^\theta$ -graded objects in ${}^H_H\mathcal{YD}$, see [AHS08, Lemma 3.2(ii)] and the discussion in [AHS08, Sect. 3.4].

Lemma 6.6. *Let $N \in \mathcal{M}_\theta$ and $i \in \mathbb{I}$. Assume that $\tilde{\Delta}(N)$ is defined.*

(1) *Let L and $(W_l)_{l \in L}$ as in Def. 6.1. Then $\mathcal{K}_i^N \simeq \otimes_{l \in L, l \neq i} \mathfrak{B}(W_l)$ as $\mathbb{N}_0^\theta$ -graded objects in ${}^H_H\mathcal{YD}$.*

(2) *Let $i, j \in \mathbb{I}$, $i \neq j$, and $n \in \mathbb{N}_0$ such that $(\text{ad}_c N_i)^n(N_j) \neq 0$. Then $\alpha_j + m\alpha_i \in \Delta(N)$ for all $0 \leq m \leq n$.*

Proof. (1) Since $\tilde{\Delta}(N)$ is defined,

$$\mathfrak{B}(V(N)) \simeq \otimes_{l \in L, l < l(i)} \mathfrak{B}(W_l) \otimes \mathfrak{B}(N_i) \otimes \otimes_{l \in L, l > l(i)} \mathfrak{B}(W_l)$$

by Lemma 4.9. Since the category ${}^H_H\mathcal{YD}$ is braided,

$$(6.12) \quad \mathfrak{B}(V(N)) \simeq \otimes_{l \in L, l \neq i} \mathfrak{B}(W_l) \otimes \mathfrak{B}(N_i).$$

Thus Eqs. (6.12), (6.11) and Lemma 4.8 imply Claim (1).

Now we prove (2). By [AHS08, Prop. 3.6] the algebra $\mathcal{K}_i^N$ is generated by the $\mathbb{N}_0^\theta$ -homogeneous subspaces $(\mathrm{ad}_c N_i)^m(N_p)$ of degree $\alpha_p + m\alpha_i$, where $p \in \mathbb{I} \setminus \{i\}$ and $m \geq 0$, and hence

$$(6.13) \quad (\mathcal{K}_i^N)_{\alpha_j + m\alpha_i} = (\mathrm{ad}_c N_i)^m(N_j).$$

On the other hand, since $\deg W_l \in (\sum_{j=1}^\theta \mathbb{N}_0 \alpha_j) \setminus \mathbb{N}_0 \alpha_i$ for all $l \in L \setminus \{l(i)\}$ by Lemma 4.9, and all tensor factors are $\mathbb{N}_0^\theta$ -graded, we obtain that

$$(6.14) \quad \left(\bigotimes_{l \in L, l \neq l(i)} \mathfrak{B}(W_l) \right)_{\alpha_j + m\alpha_i} \simeq \bigoplus_{l \in L, \deg W_l = \alpha_j + m\alpha_i} W_l.$$

Then Part (1) of the lemma and Eqs. (6.13), (6.14) give Claim (2). $\square$

The following theorem is one of the main results of [AHS08].

Theorem 6.7. [AHS08, Thm. 3.12, Lemma 3.21, Cor. 3.17] *Let $i \in \mathbb{I}$ and $N = ([N_1], \dots, [N_\theta]) \in \mathcal{M}_\theta$. Assume that N is i -finite, and let $r_i(N) = ([N'_1], \dots, [N'_\theta])$.*

(1) The family $r_i(N)$ is i -finite, $r_i^2(N) = N$, and $a_{ij}^N = a_{ij}^{r_i(N)}$ for all $j \in \mathbb{I}$. Moreover, $A^N = (a_{ij}^N)_{i,j \in \mathbb{I}}$ is a generalized Cartan matrix.

(2) Let $\deg N_j = \alpha_j$ for all $j \in \mathbb{I}$, $\deg N'_j = s_i^N(\alpha_j)$ for all $j \in \mathbb{I} \setminus \{i\}$, and $\deg N_i^ = -\alpha_i$. Then*

$$\mathfrak{B}(V(r_i(N))) \simeq \mathcal{K}_i^N \otimes \mathfrak{B}(N_i^*)$$

as $\mathbb{Z}^\theta$ -graded objects in ${}^H_H \mathcal{YD}$.

(3) The Nichols algebras $\mathfrak{B}(V(N))$ and $\mathfrak{B}(V(r_i(N)))$ have the same dimension.

Lemma 6.8. *Let $N \in \mathcal{M}_\theta$ and $i \in \mathbb{I}$. Assume that $\tilde{\Delta}(N)$ is defined and N is i -finite. Then $\tilde{\Delta}(r_i(N))$ is defined, and the map*

$$s_i^N : \tilde{\Delta}(N) \rightarrow \tilde{\Delta}(r_i(N)), \quad (C, \gamma) \mapsto (C, s_i^N(\gamma)),$$

is bijective and preserves multiplicities, that is,

$$\mathrm{mult}_N(C, \gamma) = \mathrm{mult}_{r_i(N)}(C, s_i^N(\gamma))$$

for all $(C, \gamma) \in \tilde{\Delta}(N)$.

Proof. Let $N = ([N_1], \dots, [N_\theta]) \in \mathcal{M}_\theta$ and $V(N) = N_1 \oplus \dots \oplus N_\theta$. Since $\tilde{\Delta}(N)$ is defined, there is an index set $(L, \leq)$ and a family $(W_l)_{l \in L}$ as in Def. 6.1 such that $\mathfrak{B}(V(N)) \simeq \otimes_{l \in L} \mathfrak{B}(W_l)$.

Let $r_i(N) = ([N'_1], \dots, [N'_\theta]) \in \mathcal{M}_\theta$ as in Def. 6.4, and $V(r_i(N)) = N'_1 \oplus \dots \oplus N'_\theta$. By Lemma 6.6 and Thm. 6.7(2) we obtain the decomposition

$$(6.15) \quad \mathfrak{B}(V(r_i(N))) \simeq \otimes_{l \in L, l \neq l(i)} \mathfrak{B}(W_l) \otimes \mathfrak{B}(N_i^*)$$

as $\mathbb{Z}^\theta$ -graded objects in ${}^H_H\mathcal{YD}$. We now use the group automorphism $s_i^N : \mathbb{Z}^\theta \rightarrow \mathbb{Z}^\theta$ to change the gradation. For all $l \in L$, $l \neq l(i)$, let $W'_l = W_l$ as object in ${}^H_H\mathcal{YD}$ and let $\deg(W'_l) = s_i^N(\deg(W_l))$, and $W'_{l(i)} = N_i^*$ with $\deg(N_i^*) = \alpha_i$. The isomorphism in (6.15) gives an isomorphism

$$(6.16) \quad \mathfrak{B}(V(r_i(N))) \simeq \otimes_{l \in L} \mathfrak{B}(W'_l)$$

of $\mathbb{Z}^\theta$ -graded objects in ${}^H_H\mathcal{YD}$, where $\deg N'_j = \alpha_j$ for all $j \in \mathbb{I}$. Thus

$$\begin{aligned} \tilde{\Delta}_+(r_i(N)) &= \{([W_l], s_i^N(\deg(W_l))) \mid l \in L, l \neq l(i)\} \cup \{([N_i^*], \alpha_i)\} \\ &= s_i^N \left((\tilde{\Delta}_+(N) \setminus \{([N_i], \alpha_i)\}) \cup \{([N_i^*], -\alpha_i)\} \right). \end{aligned}$$

Since $\tilde{\Delta} = \tilde{\Delta}_+ \cup \tilde{\Delta}_-$, it follows that $\tilde{\Delta}(r_i(N)) = s_i^N(\tilde{\Delta}(N))$. Moreover, for all $\gamma \in \mathbb{N}_0^\theta$, $\gamma \neq \alpha_i$, and $C \in \widehat{{}^H_H\mathcal{YD}}$,

$$\begin{aligned} \text{mult}_N(C, \gamma) &= |\{l \in L \mid C = [W_l], \gamma = \deg(W_l)\}| \\ &= |\{l \in L \mid C = [W'_l], s_i^N(\gamma) = \deg(W'_l)\}| \\ &= \text{mult}_{r_i(N)}(C, s_i^N(\gamma)), \end{aligned}$$

and $\text{mult}_N(C, \alpha_i) = 1 = \text{mult}_{r_i(N)}(C^*, \alpha_i) = \text{mult}_{r_i(N)}(C, s_i^N(\alpha_i))$ with $C = [N_i]$. This proves the lemma. $\square$

Definition 6.9. For all $M \in \mathcal{M}_\theta$ let

$$\mathcal{M}_\theta(M) = \{r_{i_1} \cdots r_{i_n}(M) \in \mathcal{M}_\theta \mid n \in \mathbb{N}_0, i_1, \dots, i_n \in \mathbb{I}\}.$$

If N is $\mathbb{I}$ -finite for all $N \in \mathcal{M}_\theta(M)$, then let

$$\mathcal{C}(M) = (\mathbb{I}, \mathcal{M}_\theta(M), (r_i)_{i \in \mathbb{I}}, (A^N)_{N \in \mathcal{M}_\theta(M)}).$$

Theorem 6.10. Let $M \in \mathcal{M}_\theta$. Assume that N is $\mathbb{I}$ -finite for all $N \in \mathcal{M}_\theta(M)$. Then $\mathcal{C}(M)$ is a Cartan scheme.

Proof. This follows from Thm. 6.7(1). $\square$

In the situation of Thm. 6.10, $\mathcal{C}(M)$ is called the *Cartan scheme* of M , and $\mathcal{W}(M) = \mathcal{W}(\mathcal{C}(M))$ is called the *Weyl groupoid* of M . It is clear by construction that $\mathcal{W}(M)$ is connected.

Theorem 6.11. *Let $M \in \mathcal{M}_\theta$. Assume that N is $\mathbb{I}$ -finite for all $N \in \mathcal{M}_\theta(M)$, and that $\tilde{\Delta}(M)$ is defined. Then*

$$\mathcal{R}(M) = (\mathcal{C}(M), (\Delta(N))_{N \in \mathcal{M}_\theta(M)})$$

is a root system of type $\mathcal{C}(M)$, called the root system of M .

If for $M \in \mathcal{M}_\theta$ the assumptions of Thm. 6.11 are satisfied, then we say that $\mathcal{R}(M)$ is defined. In this case $\tilde{\Delta}(N)$ is defined for all $N \in \mathcal{M}_\theta(M)$ by Lemma 6.8.

Proof. We have to prove (R1)–(R4) from Sect. 5.

(R1) follows from Eq. (6.6), and (R2) holds by Lemma 4.9. (R3) is a direct consequence of Lemma 6.8.

(R4) Let $w_0 = \text{id} \in \text{Aut}(\mathbb{Z}^\theta)$, and for all $n \in \mathbb{N}_0$ let

$$w_{n+1} = s_i^{r_j(r_i r_j)^n(N)} s_j^{(r_i r_j)^n(N)} w_n \in \text{Aut}(\mathbb{Z}^\theta).$$

By [HY08, Lemma 5] we obtain that

$$w_{m_{i,j;N}}(\alpha_i) = \alpha_i, \quad w_{m_{i,j;N}}(\alpha_j) = \alpha_j.$$

Note that the proof of [HY08, Lemma 5] applies, since in our case π_a in [HY08] is always the standard basis.

Let $k \in \mathbb{I} \setminus \{i, j\}$. By (R3) we have

$$w_{m_{i,j;N}}(\alpha_k) \in \alpha_k + \mathbb{Z}\alpha_i + \mathbb{Z}\alpha_j \cap \Delta((r_i r_j)^{m_{i,j;N}}(N)),$$

hence $w_{m_{i,j;N}}(\alpha_k) = \alpha_k + n_i \alpha_i + n_j \alpha_j$ for some $n_i, n_j \in \mathbb{N}_0$. Since $\alpha_k \in \Delta((r_i r_j)^{m_{i,j;N}}(N))$, it follows from (R3) that

$$\begin{aligned} w_{m_{i,j;N}}^{-1}(\alpha_k) &= w_{m_{i,j;N}}^{-1}(w_{m_{i,j;N}}(\alpha_k) - n_i \alpha_i - n_j \alpha_j) \\ &= \alpha_k - n_i \alpha_i - n_j \alpha_j \in \Delta(N). \end{aligned}$$

Hence $n_i = n_j = 0$ by (R1), that is, $w_{m_{i,j;N}} = \text{id}$. Then

$$\tilde{\Delta}(N) = w_{m_{i,j;N}}(\tilde{\Delta}(N)) = \tilde{\Delta}((r_i r_j)^{m_{i,j;N}}(N))$$

by Lemma 6.8. Thus $(r_i r_j)^{m_{i,j;N}}(N) = N$ by Lemma 6.3. $\square$

Corollary 6.12. *Let $M \in \mathcal{M}_\theta$ such that $\tilde{\Delta}(M)$ is defined and is finite. Then $\mathcal{R}(M)$ is defined, and $\mathcal{R}(M)$ is a root system of type $\mathcal{C}(M)$.*

Proof. By Thm. 6.11 it suffices to prove that $r_{i_n} \cdots r_{i_1}(M)$ is $\mathbb{I}$ -finite for all $n \in \mathbb{N}_0$ and $i_1, \dots, i_n \in \mathbb{I}$. We proceed by induction on n . The case $n = 0$ is trivial. Let $n \in \mathbb{N}_0$, $i_1, \dots, i_n \in \mathbb{I}$, and $N = r_{i_n} \cdots r_{i_1}(M)$. Since $\tilde{\Delta}(M)$ is defined and is finite, Lemma 6.8 implies that $\tilde{\Delta}(N)$ is defined and is finite. Then from Lemma 6.6(2) we obtain that for all $i, j \in \mathbb{I}$ with $i \neq j$ there exists $n_{ij} \in \mathbb{N}$ such that $(\text{ad}_c N_i)^{n_{ij}}(N_j) = 0$ in $\mathfrak{B}(V(N))$, where $N = ([N_1], \dots, [N_\theta])$. Hence N is $\mathbb{I}$ -finite. $\square$

7. FINITE ROOT SYSTEMS FOR NICHOLS ALGEBRAS

Lemma 7.1. *Let $M = ([M_1], \dots, [M_\theta]) \in \mathcal{M}_\theta$ such that $\mathcal{R}(M)$ is defined. Let $(L, \leq)$ be a totally ordered index set and $(W_l)_{l \in L}$ a family of finite-dimensional irreducible $\mathbb{Z}^\theta$ -graded objects in ${}^H_H\mathcal{YD}$, such that*

$$(7.1) \quad \mathfrak{B}(V(M)) \simeq \otimes_{l \in L} \mathfrak{B}(W_l)$$

as $\mathbb{Z}^\theta$ -graded objects in ${}^H_H\mathcal{YD}$, where $\deg(M_i) = \alpha_i$ for all $i \in \mathbb{I}$.

(1) *For all $\gamma \in \Delta_+^{\text{re}}(M)$ there is exactly one $l(\gamma) \in L$ such that $\deg W_{l(\gamma)} = \gamma$.*

(2) *If $N = ([N_1], \dots, [N_\theta]) \in \mathcal{M}_\theta(M)$, $w \in \text{Hom}(N, M)$ and $i \in \mathbb{I}$ such that $w(\alpha_i) \in \Delta_+^{\text{re}}(M)$, then $W_{l(w(\alpha_i))} \simeq N_i$.*

Proof. The decomposition (7.1) exists by assumption since $\mathcal{R}(M)$ is defined.

Let $\gamma \in \Delta_+^{\text{re}}(M)$. Then there exist $N = ([N_1], \dots, [N_\theta]) \in \mathcal{M}_\theta(M)$, $w \in \text{Hom}(N, M)$ and $i \in \mathbb{I}$ such that $w(\alpha_i) = \gamma$. Since $([N_i], \alpha_i) \in \tilde{\Delta}(N)$, Lemma 6.8 implies that $([N_i], \gamma) \in \tilde{\Delta}(M)$, that is, there exists $l \in L$ such that $\deg W_l = \gamma$ and $W_l \simeq N_i$.

Let $l, l' \in L$ with $\gamma = \deg(W_l) = \deg(W_{l'}) \in \Delta_+^{\text{re}}$. Let $N = ([N_1], \dots, [N_\theta]) \in \mathcal{M}_\theta(M)$, $w \in \text{Hom}(N, M) \subset \text{Hom}(\mathcal{W}(M))$, and $i \in \mathbb{I}$ such that $w(\alpha_i) = \gamma$. Then $w^{-1}([W_l], \gamma) = ([W_l], \alpha_i) \in \tilde{\Delta}(N)$ by Lemma 6.8, and similarly, $([W_{l'}], \alpha_i) \in \tilde{\Delta}(N)$. By Lemma 4.9 we obtain that $[W_l] = [N_i] = [W_{l'}]$. The second part of Lemma 6.8 implies that $\text{mult}_M([W_l], \gamma) = \text{mult}_N([N_i], \alpha_i) = 1$. Hence $l = l'$. This proves (1) and (2). $\square$

Theorem 7.2. *Let $M = ([M_1], \dots, [M_\theta]) \in \mathcal{M}_\theta$ such that $\mathcal{R}(M)$ is defined. Assume that $\mathcal{W}(M)$ is finite. Let $(L, \leq)$ be a totally ordered index set and $(W_l)_{l \in L}$ a family of finite-dimensional irreducible $\mathbb{Z}^\theta$ -graded objects in ${}^H_H\mathcal{YD}$, such that*

$$(7.2) \quad \mathfrak{B}(V(M)) \simeq \otimes_{l \in L} \mathfrak{B}(W_l)$$

as $\mathbb{Z}^\theta$ -graded objects in ${}^H_H\mathcal{YD}$, where $\deg(M_i) = \alpha_i$ for all $i \in \mathbb{I}$.

(1) The map $L \rightarrow \Delta_+(M)$, $l \mapsto \deg(W_l)$, is bijective. Moreover,

$$(7.3) \quad \begin{aligned} \{[W_l], [W_l^*] \mid l \in L\} &= \{[N_i] \mid i \in \mathbb{I}, \\ N &= ([N_1], \dots, [N_\theta]) \in \mathcal{M}_\theta(M)\}. \end{aligned}$$

(2) If $\gamma \in \Delta_+(M)$, then there exist $N = ([N_1], \dots, [N_\theta]) \in \mathcal{M}_\theta(M)$, $w \in \text{Hom}(N, M)$ and $i \in \mathbb{I}$ such that $\gamma = w(\alpha_i)$. In this case

$$W_{l(\gamma)} \simeq N_i,$$

where $l(\gamma)$ is the unique element in L with $\deg W_{l(\gamma)} = \gamma$.

(3) Let $i, j \in \mathbb{I}$, $i \neq j$, and $0 \leq m \leq -a_{ij}^M$. Then there is an index $l \in L$ such that $(\text{ad}_c M_i)^m(M_j) \simeq W_l$ and $\deg W_l = \alpha_j + m\alpha_i$. In particular $(\text{ad}_c M_i)^m(M_j)$ is irreducible in ${}^H_H\mathcal{YD}$.

Proof. Since $\mathcal{W}(M)$ is finite, $\Delta^{\text{re}}(M) = \Delta(M)$ by Prop. 5.2.

We first prove (1) and (2). The map $L \rightarrow \Delta_+(M)$ in (1) is surjective by definition of $\Delta_+(M)$. Hence it is bijective by Lemma 7.1(1) and the equality $\Delta^{\text{re}}(M) = \Delta(M)$. Then (2) follows from Lemma 7.1(2), since $\Delta^{\text{re}}(M) = \Delta(M)$. It remains to prove Eq. (7.3).

Let $l \in L$ and $\gamma = \deg W_l$. Then $l = l(\gamma)$ by (1), and $W_l \simeq N_i$ for some $N \in \mathcal{M}_\theta(M)$ and $i \in \mathbb{I}$ by (2). Further, if $N \in \mathcal{M}_\theta(M)$ and $i \in \mathbb{I}$ then $r_i(N) \in \mathcal{M}_\theta(M)$. Since $[r_i(N)]_i = [N_i^*]$, the right hand side of Eq. (7.3) is stable under passing to dual objects. Thus the inclusion $\subset$ holds in (3). Conversely, let $N \in \mathcal{M}_\theta(M)$ and $i \in \mathbb{I}$. Since $\mathcal{W}(M)$ is a connected groupoid, there exists $w \in \text{Hom}(N, M)$. Then Lemma 6.8 gives that $([N_i], w(\alpha_i)) \in \tilde{\Delta}(M)$, and hence $[N_i]$ is contained in the left hand side of Eq. (7.3).

Now we prove (3). Using Lemma 4.8, the isomorphism (6.11) for $N = M$ of $\mathbb{N}_0^\theta$ -graded objects in ${}^H_H\mathcal{YD}$ implies that

$$(7.4) \quad \mathcal{K}_i^M \simeq \otimes_{l \in L, l \neq l(\alpha_i)} \mathfrak{B}(W_l)$$

as $\mathbb{N}_0^\theta$ -graded objects in ${}^H_H\mathcal{YD}$. By [AHS08, Prop. 3.6] the algebra $\mathcal{K}_i^M$ is generated by the $\mathbb{N}_0^\theta$ -homogeneous subspaces $(\mathrm{ad}_c M_i)^n(M_p)$ of degree $\alpha_p + n\alpha_i$, where $p \in \mathbb{I} \setminus \{i\}$ and $n \geq 0$, and hence

$$(7.5) \quad (\mathcal{K}_i^M)_{\alpha_j + m\alpha_i} = (\mathrm{ad}_c M_i)^m(M_j).$$

On the other hand, since $\deg W_l \in (\sum_{j=1}^\theta \mathbb{N}_0 \alpha_j) \setminus \mathbb{N}_0 \alpha_i$, and all tensor factors are $\mathbb{N}_0^\theta$ -graded, we obtain that

$$(7.6) \quad \left(\bigotimes_{l \in L, l \neq i(\alpha_i)} \mathfrak{B}(W_l) \right)_{\alpha_j + m\alpha_i} \simeq W_{l(\alpha_j + m\alpha_i)}.$$

Then Eqs. (7.4), (6.13) and (6.14) give the claim of Part (3). $\square$

Theorem 7.3. *Let $M \in \mathcal{M}_\theta$. Assume that $\tilde{\Delta}(M)$ is defined. Then the following are equivalent.*

- (1) $\mathfrak{B}(V(M))$ is finite-dimensional,
- (2) (a) $\mathcal{R}(M)$ is defined and $\mathcal{W}(M)$ is finite, and
 (b) $\mathfrak{B}(N_i)$ is finite-dimensional for all $N = ([N_1], \dots, [N_\theta]) \in \mathcal{M}_\theta(M)$ and $i \in \mathbb{I}$.

Proof. (1) $\Rightarrow$ (2). Since $\mathfrak{B}(V(M))$ is finite-dimensional, N is $\mathbb{I}$ -finite for all $N \in \mathcal{M}_\theta(M)$ by Thm. 6.7(3), hence $\mathcal{R}(M)$ is defined. Further, $\Delta(M)$ is finite, and hence $\mathcal{W}(M)$ is finite by Lemma 5.1. Finally, $\dim \mathfrak{B}(V(N)) = \dim \mathfrak{B}(V(M))$ by Thm. 6.7(3), and hence $\mathfrak{B}(N_i)$ is finite-dimensional for all $i \in \mathbb{I}$ and $N = ([N_1], \dots, [N_\theta]) \in \mathcal{M}_\theta(M)$.

(2) $\Rightarrow$ (1). From (2a) and Lemma 5.1 we obtain that $\Delta(M)$ is finite. Hence $\mathfrak{B}(V(M))$ is finite-dimensional by (2b) and Thm. 7.2 (1) and (2). $\square$

Recall that $\tilde{\Delta}(M)$ is defined in particular if all tensor powers of $V(M)$ are semisimple.

Let $M \in \mathcal{M}_\theta$. Then M is called *standard*, see [AHS08, Def. 3.23], if N is $\mathbb{I}$ -finite and $A^N = A^M$ for all $N \in \mathcal{M}_\theta(M)$.

Corollary 7.4. *Let $M \in \mathcal{M}_\theta$. Assume that $\tilde{\Delta}(M)$ is defined, and that M is standard. Then $\mathcal{R}(M)$ is defined, and the following are equivalent.*

- (1) $\mathfrak{B}(V(M))$ is finite-dimensional,
- (2) (a) A^M is a Cartan matrix of finite type,

- (b) $\mathfrak{B}(N_i)$ is finite-dimensional for all $N = ([N_1], \dots, [N_\theta]) \in \mathcal{M}_\theta(M)$ and $i \in \mathbb{I}$.

Proof. This follows from Thm. 7.3 and Cor. 5.4. $\square$

8. APPLICATIONS FOR FINITE GROUPS

In this section let $\mathbb{k}$ be an algebraically closed field of characteristic 0, and let $H = \mathbb{k}G$ be the group algebra of a finite group G . For any $g \in G$ let $\mathcal{O}_g$ denote the conjugacy class of g and G^g the centralizer of g in G . Let $\triangleright$ denote the adjoint action in G , that is, $g \triangleright h = ghg^{-1}$ for all $g, h \in G$. The category ${}^H_H\mathcal{YD}$ will be denoted by ${}^G_G\mathcal{YD}$.

Let $g \in G$ and M an irreducible G^g -module. Then $kG \otimes_{kG^g} M$ is an irreducible object in ${}^G_G\mathcal{YD}$, where

$$h \cdot (h' \otimes m) = hh' \otimes m, \quad \delta(h \otimes m) = hgh^{-1} \otimes (h \otimes m)$$

for all $h, h' \in G$, $m \in M$. Any irreducible object in ${}^G_G\mathcal{YD}$ arises in this way. The Yetter-Drinfeld module $V = kG \otimes_{kG^g} M$ can be written as

$$V = \oplus_{s \in \mathcal{O}_g} V_s, \quad V_s = \{v \in V \mid \delta(v) = s \otimes v\},$$

and $V_g = 1 \otimes M$. For all $s \in \mathcal{O}_g$, V_s is an irreducible G^s -module, and $h \cdot V_s = V_{h \triangleright s}$ for all $h \in G$. There exists $q_V \in \mathbb{k}^*$ such that $s \cdot v = q_V v$ for all $v \in V_s$, $s \in \mathcal{O}_g$.

Let $g, h \in G$. We say that $\mathcal{O}_g$ and $\mathcal{O}_h$ commute, if $st = ts$ for all $s \in \mathcal{O}_g$, $t \in \mathcal{O}_h$.

Proposition 8.1. *Let $g, h \in G$ and $V = \oplus_{s \in \mathcal{O}_g} V_s$, $W = \oplus_{t \in \mathcal{O}_h} W_t$ be irreducible objects in ${}^G_G\mathcal{YD}$.*

(1) *If $(\text{ad}_c V)(W) = 0$ in $\mathfrak{B}(V \oplus W)$, then $\mathcal{O}_g$ and $\mathcal{O}_h$ commute.*

(2) *If $(\text{ad}_c V)^2(W) = 0$ in $\mathfrak{B}(V \oplus W)$, then $\mathcal{O}_g$ commutes with itself or with $\mathcal{O}_h$.*

Proof. (1) By Prop. 6.5, $(\text{ad}_c V)(W)$ is isomorphic to $(\text{id} - c^2)(V \otimes W)$ in ${}^G_G\mathcal{YD}$. If $s \in \mathcal{O}_g$, $t \in \mathcal{O}_h$, $st \neq ts$, then

$$c^2(V_s \otimes W_t) = V_{st \triangleright s} \otimes W_{s \triangleright t},$$

and hence $(\text{id} - c^2)(V_s \otimes W_t) \neq 0$.

(2) It suffices to show that if $(\text{ad}_c V)^2(W) = 0$ and $gh \neq hg$, then $rg = gr$ for all $r \in \mathcal{O}_g \setminus \{g\}$. We note that if $r \in \mathcal{O}_g$, $v \in V_r$, $v' \in V_g$, $w \in W_h$, $v \otimes v' \otimes w \neq 0$, then $(S_2 \otimes \text{id})T_2(v \otimes v' \otimes w)$ is the sum of eight nonzero homogeneous terms of degrees

$$\begin{aligned} & (r, g, h), (r \triangleright g, r, h), (r, gh \triangleright g, g \triangleright h), (rgh \triangleright g, r, g \triangleright h), \\ & (r \triangleright g, rh \triangleright r, r \triangleright h), (rgh \triangleright r, r \triangleright g, r \triangleright h), \\ & (rgh \triangleright g, rghg^{-1} \triangleright r, rg \triangleright h), (rgh \triangleright r, rgh \triangleright g, rg \triangleright h). \end{aligned}$$

On the other hand, $(S_2 \otimes \text{id})T_2(v \otimes v' \otimes w) = 0$ by Prop. 6.5 and the assumption $(\text{ad}_c V_r)(\text{ad}_c V_g)(W_h) = 0$. The vanishing of the sum of these eight terms gives information about the corresponding degrees.

Let first $r \in \mathcal{O}_g$ such that $rh \neq hr$ and $r \neq g$. By comparing degrees we get $rg \triangleright h = h$ and $rgh \triangleright g = r$. Hence $r = gh \triangleright g$ is uniquely determined. Since $h \triangleright g$ and $h^{-1} \triangleright g$ do not commute with h and are different from g , we get

$$(8.1) \quad r = h^{-1} \triangleright g = h \triangleright g = gh \triangleright g.$$

Thus $g = h^{-1}gh \triangleright g = r \triangleright g$ by the last and first equations in Eq. (8.1), that is, $rg = gr$.

Let now $r \in \mathcal{O}_g$ with $rh = hr$. Then $r \neq g$, and by comparing degrees one gets $rg \triangleright h = g \triangleright h$ and $r \triangleright g = g$. This proves the claim. $\square$

Let $\mathcal{F}(G)$ be the set of all conjugacy classes $\mathcal{O}$ of G such that $\dim \mathfrak{B}(V) < \infty$ for some $V = \oplus_{s \in \mathcal{O}} V_s \in {}^G\mathcal{YD}$.

Theorem 8.2. *Assume that any two conjugacy classes in $\mathcal{F}(G)$ do not commute. Let $U \in {}^G\mathcal{YD}$. If $\mathfrak{B}(U)$ is finite-dimensional, then U is irreducible in ${}^G\mathcal{YD}$.*

Proof. Recall that the category ${}^G\mathcal{YD}$ is semisimple, and that embeddings of Yetter-Drinfeld modules induce embeddings of the corresponding Nichols algebras. Hence it suffices to prove that $\mathfrak{B}(V \oplus W)$ is infinite dimensional for all irreducible objects $V, W \in {}^G\mathcal{YD}$.

Let $g, h \in G$, $V = \oplus_{s \in \mathcal{O}_g} V_s$, $W = \oplus_{t \in \mathcal{O}_h} V_t$ irreducible objects in ${}^G\mathcal{YD}$ and $M = ([V], [W]) \in \mathcal{M}_2$. Assume that $\dim \mathfrak{B}(V \oplus W) < \infty$. Then $\dim \mathfrak{B}(V), \dim \mathfrak{B}(W) < \infty$, and hence $\mathcal{O}_g, \mathcal{O}_h \in \mathcal{F}(G)$.

If $a_{12}^M = 0$, then $(\text{ad}_c V)(W) = 0$, and hence Prop. 8.1(1) gives that $\mathcal{O}_g$ and $\mathcal{O}_h$ commute. If $a_{12}^M = -1$, then $(\text{ad}_c V)^2(W) = 0$, and Prop. 8.1(2) gives that $\mathcal{O}_g$ commutes with $\mathcal{O}_g$ or with $\mathcal{O}_h$. Thus the assumption in the theorem yields that $a_{12}^M < -1$. Similarly, $a_{21}^M < -1$. By Thm. 6.7(3), $\dim \mathfrak{B}(N_1 \oplus N_2) = \dim \mathfrak{B}(M_1 \oplus M_2)$ for all $N \in \mathcal{M}_2(M)$, $N = ([N_1], [N_2])$, and hence the above arguments give that $a_{12}^N, a_{21}^N < -1$ for all $N \in \mathcal{M}_2(M)$. Since $\text{Hom}(\mathcal{W}(M))$ is finite by Thm. 7.3, the theorem follows from Prop. 5.7. $\square$

Recall that if $V = V_1$, then $\dim \mathfrak{B}(V) = \infty$, since the characteristic of $\mathbb{k}$ is 0. Thus $\mathcal{O}_1 \notin \mathcal{F}(G)$.

Corollary 8.3. *Let G be a nonabelian simple group. Let $U \in {}^G\mathcal{YD}$. If $\mathfrak{B}(U)$ is finite-dimensional, then U is irreducible in ${}^G\mathcal{YD}$.*

Proof. Let $\mathcal{O}', \mathcal{O}''$ be two commuting conjugacy classes of G . Then the subgroups $G' = \langle \mathcal{O}' \rangle$, $G'' = \langle \mathcal{O}'' \rangle$ of G are normal. Since G is simple, G', G'' are either 1 or G . Since $\mathcal{O}', \mathcal{O}''$ commute, $[G', G''] = 1$. Hence $G' = 1$ or $G'' = 1$, and hence $\mathcal{O}' = 1$ or $\mathcal{O}'' = 1$. But $1 \notin \mathcal{F}(G)$, and the claim follows from Thm. 8.2. $\square$

Corollary 8.4. *Let $n \in \mathbb{N}$, $n \geq 3$, and assume that $G = \mathbb{S}_n$ is the symmetric group. Let $U \in {}^G\mathcal{YD}$. If $\mathfrak{B}(U)$ is finite-dimensional, then U is irreducible in ${}^G\mathcal{YD}$.*

Proof. For $n \geq 5$ the group $\mathbb{S}_n$ has a simple group of index two. Along the lines of the proof of Cor. 8.3 it is easy to show that $\mathbb{S}_n$, $n \geq 5$, does not possess commuting nontrivial (i.e. different from $\{1\}$) conjugacy classes. In the usual cycle notation, the only pairs of commuting nontrivial conjugacy classes of $\mathbb{S}_n$ are $(\mathcal{O}_{(123)}, \mathcal{O}_{(123)})$ for $n = 3$ and $(\mathcal{O}_{(12)(34)}, \mathcal{O}_{(12)(34)})$ for $n = 4$. By [AZ07, Thm. 1], $\mathcal{O}_{(123)} \notin \mathcal{F}(\mathbb{S}_3)$ and $\mathcal{O}_{(12)(34)} \notin \mathcal{F}(\mathbb{S}_4)$. Thus the claim follows from Thm. 8.2. $\square$

For the dihedral groups $\mathbb{D}_n$, n odd, an alternative proof of [AHS08, Thm. 4.8] can be given as another application of Thm. 8.2.

We continue with some other consequences of our theory which hold for all finite groups.

Proposition 8.5. *Let $g, h \in G$ and $V = \oplus_{s \in \mathcal{O}_g} V_s$, $W = \oplus_{t \in \mathcal{O}_h} W_t$ be irreducible objects in ${}^G\mathcal{YD}$. Assume that $(\text{ad}_c V)(W)$ is irreducible.*

(1) *$stst = tsts$ for all $s \in \mathcal{O}_g$, $t \in \mathcal{O}_h$.*

(2) *There is at most one double coset $G^h x G^g$ in $G^h \backslash G / G^g$, $x \in G$, such that $(x \triangleright g)h \neq h(x \triangleright g)$. If it exists, then $(\text{ad}_c V)(W) \neq 0$ and $q_{(\text{ad}_c V)(W)} = -q_V q_W$.*

Proof. By Prop. 6.5, $(\text{ad}_c V)(W)$ is isomorphic to $(\text{id} - c^2)(V \otimes W)$ in ${}^G\mathcal{YD}$. Let $s \in \mathcal{O}_g$, $t \in \mathcal{O}_h$, $v \in V_s$, and $w \in W_t$. Then

$$c^2(v \otimes w) = c((s \cdot w) \otimes v) = sts^{-1} \cdot v \otimes s \cdot w.$$

Since $s \cdot v = q_V v$ and $t \cdot w = q_W w$, we obtain that

$$(8.2) \quad (\text{id} - c^2)(v \otimes w) = v \otimes w - q_V^{-1} q_W^{-1} st \cdot (v \otimes w).$$

Since $(\text{ad}_c V)(W)$ is irreducible, there exists $q \in \mathbb{k}^*$ such that

$$(8.3) \quad st \cdot (\text{id} - c^2)(v \otimes w) = q(\text{id} - c^2)(v \otimes w).$$

Comparing degrees we obtain that $st \triangleright t = t$ or $(st)^2 \triangleright t = t$. Thus $(st)^2 = (ts)^2$. This proves (1).

If $X \subset G$ is a set of double coset representatives for $G^h \backslash G / G^g$, then

$$V \otimes W = \oplus_{x \in X} \mathbb{k}G \cdot ((x \cdot V_g) \otimes W_h)$$

is a decomposition into Yetter-Drinfeld modules over G . By Eq. (8.2),

$$(\text{id} - c^2)(V \otimes W) = \oplus_{x \in X} \mathbb{k}G \cdot (\text{id} - c^2)((x \cdot V_g) \otimes W_h)$$

is a decomposition into Yetter-Drinfeld modules over G . Let $x \in X$. Assume that $(x \triangleright g)h \neq h(x \triangleright g)$. Then $(\text{id} - c^2)((x \cdot V_g) \otimes W_h) \neq 0$ by Eq. (8.2). Since $(\text{ad}_c V)(W)$ is irreducible by assumption, Prop. 6.5 implies that $(\text{id} - c^2)(V \otimes W)$ is irreducible. Thus $(\text{id} - c^2)((y \cdot V_g) \otimes W_h) = 0$ and hence $(y \triangleright g)h = h(y \triangleright g)$ for all $y \in X \setminus \{x\}$. The claim $q_{(\text{ad}_c V)(W)} = -q_V q_W$ follows from Eq. (8.3) for $s = x \triangleright g$, $t = h$. $\square$

Theorem 8.6. *Let $g, h \in G$ and $V = \oplus_{s \in \mathcal{O}_g} V_s$, $W = \oplus_{t \in \mathcal{O}_h} W_t$ be irreducible objects in ${}^G\mathcal{YD}$. If $\mathfrak{B}(V \oplus W)$ is finite-dimensional, then the following hold.*

(1) *For all $s \in \mathcal{O}_g$ and $t \in \mathcal{O}_h$, $(st)^2 = (ts)^2$.*

(2) There is at most one double coset $G^h x G^g$ in $G^h \backslash G / G^g$, $x \in G$, such that $(x \triangleright g)h \neq h(x \triangleright g)$.

(3) For all $s \in \mathcal{O}_g$, $t \in \mathcal{O}_h$ with $st \neq ts$ there is an irreducible object $U = \bigoplus_{r \in \mathcal{O}_{st}} U_r \in {}_G^G \mathcal{YD}$ satisfying $q_U = -q_V q_W$ and $\dim \mathfrak{B}(U) < \infty$.

Proof. By Thms. 7.3 and 7.2(3), $(\text{ad}_c V)(W)$ is either 0 or irreducible. Then (1) and (2) follow from Prop. 8.5(2).

(3) Assume that $s \in \mathcal{O}_g$, $t \in \mathcal{O}_h$ such that $st \neq ts$. Then $U = (\text{ad}_c V_s)(W_t) \neq 0$ by Eq. (8.2), hence $U = \bigoplus_{r \in \mathcal{O}_{st}} U_r$ is irreducible. Therefore $q_U = -q_V q_W$ by Prop. 8.5(2), and $\mathfrak{B}(U)$ is finite-dimensional by Thm. 7.2 and since $\mathfrak{B}(V \oplus W)$ is finite-dimensional. $\square$

REFERENCES

- [AF08] N. Andruskiewitsch and F. Fantino, *New techniques for pointed Hopf algebras*, Preprint arXiv:0803.3486 (2008), 29 pages.
- [AHS08] N. Andruskiewitsch, I. Heckenberger, and H.-J. Schneider, *The Nichols algebra of a semisimple Yetter-Drinfeld module*, Preprint arXiv:0803.2430 (2008), 52 pages.
- [AS98] N. Andruskiewitsch and H.-J. Schneider, *Lifting of quantum linear spaces and pointed Hopf algebras of order p^3* , J. Algebra **209** (1998), 658–691.
- [AS02a] ———, *Finite quantum groups over abelian groups of prime exponent*, Ann. Scient. Éc. Norm. Sup. **35** (2002), 1–26.
- [AS02b] ———, *Pointed Hopf algebras*, New Directions in Hopf Algebras, MSRI Publications, vol. 43, Cambridge University Press, 2002.
- [AS05] ———, *On the classification of finite-dimensional pointed Hopf algebras*, Accepted for publication in Ann. Math., Preprint math.QA/0502157 (2005), 43 pages.
- [AZ07] N. Andruskiewitsch and S. Zhang, *On pointed Hopf algebras associated to some conjugacy classes in S_n* , Proc. Amer. Math. Soc. (2007), 2723–2731.
- [Bou68] N. Bourbaki, *Groupes et algèbres de Lie, ch. 4, 5 et 6*, Éléments de mathématique, Hermann, Paris, 1968.
- [CH08] M. Cuntz and I. Heckenberger, *Weyl groupoids with at most three objects*, Preprint (2008), 31 pages.
- [GH07] M. Graña and I. Heckenberger, *On a factorization of graded Hopf algebras using Lyndon words*, J. Algebra **314** (2007), 324–343.
- [Hec06a] I. Heckenberger, *Classification of arithmetic root systems*, Preprint math.QA/0605795 (2006), 67 pages.
- [Hec06b] ———, *The Weyl groupoid of a Nichols algebra of diagonal type*, Invent. Math. **164** (2006), 175–188.

- [Hec08] ———, *Rank 2 Nichols algebras with finite arithmetic root system*, Algebr. Represent. Theory **11** (2008), 115–132.
- [HY08] I. Heckenberger and H. Yamane, *A generalization of Coxeter groups, root systems, and Matsumoto’s theorem*, Math. Z. **259** (2008), 255–276.
- [Kac90] V.G. Kac, *Infinite dimensional Lie algebras*, Cambridge Univ. Press, 1990.
- [Kha99] V. Kharchenko, *A quantum analogue of the Poincaré–Birkhoff–Witt theorem*, Algebra and Logic **38** (1999), no. 4, 259–276.
- [Lus93] G. Lusztig, *Introduction to quantum groups*, Birkhäuser, Boston, MA, 1993.
- [Maj95] S. Majid, *Foundations of quantum group theory*, Cambridge University Press, 1995.
- [Mon93] S. Montgomery, *Hopf algebras and their actions on rings*, Amer. Math. Soc., Providence, RI, 1993.
- [Sch96] P. Schauenburg, *A characterization of the borel-like subalgebras of quantum enveloping algebras*, Comm. Algebra **24** (1996), 2811–2823.
- [Wor89] S.L. Woronowicz, *Differential calculus on compact matrix pseudogroups (quantum groups)*, Comm. Math. Phys. **122** (1989), no. 1, 125–170.

ISTVÁN HECKENBERGER, MATHEMATISCHES INSTITUT, UNIVERSITÄT MÜNCHEN,
THERESIENSTR. 39, D-80333 MUNICH, GERMANY

E-mail address: i.heckenberger@googlemail.com

HANS-JÜRGEN SCHNEIDER, MATHEMATISCHES INSTITUT, UNIVERSITÄT MÜNCHEN,
THERESIENSTR. 39, D-80333 MUNICH, GERMANY

E-mail address: Hans-Juergen.Schneider@mathematik.uni-muenchen.de